CANADIAN THESES ON MICROFICHE

I.S.B.N.

THESES CANADIENNES SUR MICROFICHE

National Library of Canada
Collections Development Branch

Canadian Theses on
Microfiche Service

Ottawa, Canada
K1A 0N4

Bibliothèque nationale du Canada
Direction du développement des collections

Service des thèses canadiennes
sur microfiche

NOTICE

The quality of this microfiche is heavily dependent upon the quality of the original thesis submitted for microfilming. Every effort has been made to ensure the highest quality of reproduction possible.

If pages are missing, contact the university which granted the degree.

Some pages may have indistinct print especially if the original pages were typed with a poor typewriter ribbon or if the university sent us a poor photocopy.

Previously copyrighted materials (journal articles, published tests, etc.) are not filmed.

Reproduction in full or in part of this film is governed by the Canadian Copyright Act, R.S.C. 1970, c. C-30. Please read the authorization forms which accompany this thesis.

THIS DISSERTATION
HAS BEEN MICROFILMED
EXACTLY AS RECEIVED

AVIS

La qualité de cette microfiche dépend grandement de la qualité de la thèse soumise au microfilmage. Nous avons tout fait pour assurer une qualité supérieure de reproduction.

S'il manque des pages, veuillez communiquer avec l'université qui a conféré le grade.

La qualité d'impression de certaines pages peut laisser à désirer, surtout si les pages originales ont été dactylographiées à l'aide d'un ruban usé ou si l'université nous a fait parvenir une photocopie de mauvaise qualité.

Les documents qui font déjà l'objet d'un droit d'auteur (articles de revue, examens publiés, etc.) ne sont pas microfilmés.

La reproduction, même partielle, de ce microfilm est soumise à la Loi canadienne sur le droit d'auteur, SRC 1970, c. C-30. Veuillez prendre connaissance des formules d'autorisation qui accompagnent cette thèse.

LA THÈSE A ÉTÉ
MICROFILMÉE TELLE QUE
NOUS L'AVONS REÇUE

NL-339 (r. 82/08)

Canada

0-315-09037-5

National Library of Canada | Bibliothèque nationale du Canada

Canadian Theses Division | Division des thèses canadiennes

154 Ottawa, Canada K1A 0N4

56954

PERMISSION TO MICROFILM — AUTORISATION DE MICROFILMER

• Please print or type — Écrire en lettres moulées ou dactylographier

Full Name of Author — Nom complet de l'auteur

Andrew Allan Turner

Date of Birth — Date de naissance	Country of Birth — Lieu de naissance
Nov. 23, 1949.	Canada

Permanent Address — Résidence fixe

613 Third Avenue South
Kenora, Ontario
Canada P9N 1H4

Title of Thesis — Titre de la thèse

Effects of Different Training Methods On Flexibility

University — Université

University of Alberta

Degree for which thesis was presented — Grade pour lequel cette thèse fut présentée

Ph.D.

Year this degree conferred — Année d'obtention de ce grade	Name of Supervisor — Nom du directeur de thèse
1982	Dr. M Singh

Permission is hereby granted to the NATIONAL LIBRARY OF CANADA to microfilm this thesis and to lend or sell copies of the film.

The author reserves other publication rights, and neither the thesis nor extensive extracts from it may be printed or otherwise reproduced without the author's written permission.

L'autorisation est, par la présente, accordée à la BIBLIOTHÈQUE NATIONALE DU CANADA de microfilmer cette thèse et de prêter ou de vendre des exemplaires du film.

L'auteur se réserve les autres droits de publication; ni la thèse ni de longs extraits de celle-ci ne doivent être imprimés ou autrement reproduits sans l'autorisation écrite de l'auteur.

Date	Signature
April 23, 1982	A. Allan Turner

NL-91 (4/77)

THE UNIVERSITY OF ALBERTA

EFFECTS OF DIFFERENT TRAINING METHODS ON FLEXIBILITY

BY

(C)

ANDREW ALLAN TURNER

A THESIS

SUBMITTED TO THE FACULTY OF GRADUATE STUDIES AND RESEARCH

IN PARTIAL FULFILLMENT OF THE REQUIREMENTS FOR THE DEGREE

OF DOCTOR OF PHILOSOPHY

DEPARTMENT OF PHYSICAL EDUCATION

EDMONTON, ALBERTA

SPRING, 1982

THE UNIVERSITY OF ALBERTA

RELEASE FORM

NAME OF AUTHOR ANDREW ALLAN TURNER

TITLE OF THESIS EFFECTS OF DIFFERENT TRAINING METHODS ON FLEXIBILITY

DEGREE FOR WHICH THESIS WAS PRESENTED DOCTOR OF PHILOSOPHY

YEAR THIS DEGREE GRANTED 1982

Permission is hereby granted to THE UNIVERSITY OF ALBERTA LIBRARY to reproduce single copies of this thesis and to lend or sell such copies for private, scholarly or scientific research purposes only.

The author reserves other publication rights, and neither the thesis nor extensive extracts from it may be printed or otherwise reproduced without the author's written permission.

(Signed) A. Allan Turner

PERMANENT ADDRESS:
613 Third Avenue South
Kenora, Ontario
Canada P9N 1Y4

DATED 82.04.23 1982

THE UNIVERSITY OF ALBERTA

FACULTY OF GRADUATE STUDIES AND RESEARCH

The undersigned certify that they have read, and recommend to the Faculty of Graduate Studies and Research, for acceptance, a thesis entitled EFFECTS OF DIFFERENT TRAINING METHODS ON FLEXIBILITY

submitted by ANDREW ALLAN TURNER

in partial fulfillment of the requirements for the degree of Doctor of Philosophy in Physical Education.

Mohan Singh
Supervisor

S. W. Mendryk

Magnus

Skumar

David Wyse

K. D. Coutts
External Examiner

Date 82.04.19

DEDICATION

To my wife, my mother, and my father,
for all their assistance, support,
and love during my years as a student.

ABSTRACT

The purpose of this study was to investigate the effects of six different standardized stretch training methods on active flexibility for shoulder extension and ankle plantar flexion. The different training methods were: 1) 3S (concentric, isometric and concentric contractions with external force), 2) 2S (passive extension), 3) SS (concentric and isometric contractions), 4) ES (eccentric contraction and passive extension), 5) BS (concentric contraction), and 6) CS (concentric, isometric and concentric contractions). Forty-two female volunteer students ranging in age from 12 to 14 years were selected as subjects. They were tested at the beginning of the study (pre-test), after two weeks, four weeks, and six weeks (post-test), and after the two week retention period. The modified Leighton Flexometer was utilized to test active range of motion for the ankle and shoulder joints.

Data was analyzed using two three-way analyses of variance (Greenhouse-Geisser test on repeated measures) and post-hoc Scheffé test ($\alpha = 0.05$). Analysis of data on ankle joints revealed that 3S, 2S, ES, and CS training method increased within-session flexibility when compared to the control ($p < 0.05$). In addition, 3S training was superior to CS training in relation to within-session gains ($p < 0.05$). The data on shoulder joints showed that only the 3S training method increased within-session flexibility when compared to the control group. Also, the 3S training method was

the only one which consistently increased range of motion on both the ankle and shoulder joints for within-session training.

ACKNOWLEDGEMENTS

I wish to express my appreciation to:

1) Dr. M. Singh, my advisor, for his encouragement and guidance;

2) Dr. T. Maguire, Dr. S. Mendryk, Dr. S. Kumar and Dr. D. Magee, my committee members, for their scholarly assistance and co-operation;

3) Dr. K. D. Coutts, my external examiner, for his insight and constructive criticism;

4) Mr. R. Harvey, principal of Lakewood Intermediate School, Kenora, Ontario;

5) Marcel Bouffard, my close friend and colleague, for sharing his time and knowledge;

6) Chet Atkins, for his photography;

7) Cheryl Luchkow, for her typing; and

8) Cheryl Turner, my wife, for proofreading my thesis.

TABLE OF CONTENTS

CHAPTER		PAGE
I	STATEMENT OF THE PROBLEM	1
	Introduction	1
	Problem	4
	Limitations of the Study	5
	Delimitations of the Study	6
	Definitions of the Study	6
II	REVIEW OF LITERATURE	10
	Overview of Flexibility	10
	Flexibility Training	10
	Summary	22
III	METHODOLOGY	26
	Subjects	26
	Randomization	26
	Selection	26
	Assignment	26
	Modification	27
	Experimental Design	27
	Testing Apparatus	29
	Modified Leighton Flexometer	31
	Apparatus	33
	Testing Procedures	33
	Flexibility Tests	35
	Training Apparatus	36
	Training Procedures	37
	Flexibility Training	38
	Scientific Stretch for Sport Training (3S)	38

CHAPTER PAGE

Static Stretch Training (2S) 39

Slow Active Stretch Training (SS) . . . 39

Eccentric Stretch Training (ES) 39

Ballistic Stretch Training (BS) 40

Contract Relax Stretch Training (CS) . . 40

Summary of Test and Training Format 41

Pre-test Orientation 41

Initial Test 41

Training Program 41

Training Tests 41

Post-training Period 43

Post-training Test 43

IV RESULTS AND DISCUSSION 45

Ankle Plantar Flexion 46

Shoulder Extension 47

Discussion of Treatment Over Time 61

Joint Differences for Within-Session Training 63

Scientific Investigations on 3S 64

Theoretical Explanation 67

Future Research 72

Summary 72

V SUMMARY, CONCLUSIONS AND RECOMMENDATIONS . . . 73

Summary 73

Conclusions 74

Ankle Plantar Flexion 74

Shoulder Extension 74

Comparison Across Joints 74

CHAPTER PAGE

Recommendations 75

BIBLIOGRAPHY 76

APPENDIX A. FLEXIBILITY TRAINING METHODS 83

Scientific Stretch for Sport Training 84

Static Stretch Training 95

Slow Active Stretch Training 104

Eccentric Stretch Training 111

Ballistic Stretch Training 122

Contract Relax Stretch Training . . . 129

APPENDIX B. FLEXIBILITY TESTS AND APPARATUS 140

Flexibility Tests 141

Shoulder Extension Test 141

Ankle Plantar Flexion Test 145

Apparatus 149

APPENDIX C. CORRESPONDENCE 151

APPENDIX D. RAW DATA 155

Ankle Data 156

Shoulder Data 158

APPENDIX E. TABLES OF MEANS 160

Ankle Data 161

Shoulder Data 165

LIST OF TABLES

Table	Description	Page
1	Effects of Different Stretch Training Methods on Flexibility	23
2	Summary of Analysis of Variance on the Ankle Data	48
3	Scheffé Post-Hoc Contrasts Among Levels of Factor B (Week) on the Ankle Data	50
4	Scheffé Post-Hoc Contrasts Among Levels of Factor C (Session) for Given Levels of Factor A (Treatment) on the Ankle Data	52
5	Scheffé Post-Hoc Contrasts Among Levels of Factor A (Treatment) for Significant C_1 and C_2 Differences on the Ankle Data	54
6	Summary of Analysis of Variance on the Shoulder Data	55
7	Scheffé Post-Hoc Contrasts Among Levels of Factor B (Week) on the Shoulder Data	57
8	Scheffé Post-Hoc Contrasts Among Levels of Factor C (Session) for Given Levels of Factor A (Treatment) on the Shoulder Data	59
9	Neural Mechanisms Underlying Training Methods	71
A	Raw Scores (in Degrees) of Ankle Joint Flexibility for Each Subject	156
B	Raw Scores (in Degrees) of Ankle Joint Flexibility for Each Subject	157
C	Raw Scores (in Degrees) of Shoulder Joint Flexibility for Each Subject	158
D	Raw Scores (in Degrees) of Shoulder Joint Flexibility for Each Subject	159
E	Treatment(A) by Week(B) by Session (C) Summary Table of Cell Means for Ankle Data	161
F	Treatment(A) by Week(B) Summary Table of Cell Means for Ankle Data	162

Table	Description	Page
G	Treatment(A) by Session(C) Summary Table of Cell Means for Ankle Data	163
H	Week(B) by Session(C) Summary Table of Cell Means for Ankle Data	164
I	Treatment(A) by Week(B) by Session(C) Summary Table of Cell Means for Shoulder Data	165
J	Treatment(A) by Week(B) Summary Table of Cell Means for Shoulder Data	166
K	Treatment(A) by Session(C) Summary Table of Cell Means for Shoulder Data	167
L	Week(B) by Session(C) Summary Table of Cell Means for Shoulder Data	168

LIST OF FIGURES

Figure		Page
1	Subject A: Active Flexibility	2
2	Subject B: Passive Flexibility	2
3	Design of the study	28
4	Front view of the modified Leighton Flexometer	32
5	Back view of the modified Leighton Flexometer	32
6	Shoulder extension table	34
7	Ankle plantar flexion table	34
8	Shoulder extension test	36
9	Ankle plantar flexion test	36
10	Subject is flexed at knee	42
11	Partner is supported by feet	42
12	Schematic diagram of testing and training format	44
13	Mean flexibility scores for each treatment, week, and session on the ankle joint	49
14	Mean flexibility scores for each week on the ankle joint	51
15	Mean flexibility scores within-session for each treatment on the ankle joint	53
16	Mean flexibility scores for each treatment, week, and session on the shoulder joint	56
17	Mean flexibility scores for each week on the shoulder joint	58
18	Mean flexibility scores within-session for each treatment on the shoulder joint	60
A	Test cylinder for ankle plantar flexion test	149
B	Test strap for shoulder extension test	149
C	The Leighton Flexometer	150

CHAPTER I

STATEMENT OF THE PROBLEM

Introduction

Flexibility is defined as "the range of possible movement in a joint (as in the hip joint) or series of joints (as when the spinal column is involved)" (de Vries, 1976, p. 432). This range of motion can further be divided into two phases - the active and the passive. The former refers to the range of motion produced by a voluntary contraction of the muscles, and the latter to the range of motion produced by use of force from outside the joint (e.g., force from a partner) (Hartley, 1976). The distinction is illustrated in Figure 1, which shows Subject A demonstrating active flexibility and in Figure 2, Subject B demonstrating passive flexibility for shoulder joint extension.

Consequently, flexibility can be defined as the degree to which a joint(s) is free to move through its range of motion.

The importance of flexibility was stated by Hartley (1976), listing six aspects of physical activity where flexibility plays a crucial role in enhancing:

1) Injury prevention,
2) Shock absorption,
3) Balance and agility,
4) Aesthetic impression,



Figure 1. Subject A: Active Flexibility



Figure 2. Subject B: Passive Flexibility

5) Performance efficiency, and

6) Joint amplitude.

Flexibility has been acknowledged only recently by both the practitioner and researcher to be a centrally important human performance variable (Corbin and Noble, 1980). Schultz (1979) stated that flexibility, for years the overlooked sibling of strength, endurance, and speed, has come to be appreciated for its own virtues. Jesse (1979) drew attention to the misuse of strength training with regards to the loss of specific joint flexibility (e.g., ankle flexibility). A few years earlier, Cooper and Fair (1977) called for stretching exercises to be brought out from under the piles of weight-lifting machine literature, taught properly, and implemented into every athlete's year-round training program.

From a coaching perspective, Hogg (1977) in a comprehensive review of the literature on flexibility training concluded:

> In the area of flexibility for athletes there are few valuable studies. Comparative research into the best stretching methods has been difficult because of the presence of many uncontrolled variables and the limited availability or normative data (p. 37).

From a researcher's perspective, Schultz (1979) stated that de Vries is recognized as one of the foremost investigators on the nature and effects of flexibility training. He finds it unfortunate there is no large body of evidence from other laboratory and clinical studies to support, or reject, the use of specific stretch training methods (e.g., static stretching). From a therapist's perspective, Cherry (1980)

called for research to establish which methods are most effective, development of new techniques and a scientific rationale of the most effective methods.

The review of the literature revealed that there were several popular flexibility training methods, each purporting to cause significant improvements in joint range of motion. Scientific Stretching for Sport (i.e., 3S) required a two second concentric contraction, a six second isometric contraction and a two second concentric contraction, with light force from a partner (Holt, 1973). Static Stretching (i.e., 2S) consisted of a two second passive extension and an eight second held passive extension (de Vries, 1962). Slow Active Stretching (i.e., SS) involved a two second concentric contraction and an eight second isometric contraction (Jacobs, 1976). Eccentric Stretching (i.e., ES) entailed a five second eccentric contraction and a five second passive extension. Ballistic Stretching (i.e., BS) required a one second concentric contraction (McCue, 1952). Contract Relax Stretching (i.e., CS) consisted of a two second concentric contraction, a six second isometric contraction and a two second concentric contraction (Moore and Hutton, 1980) (see Appendix A).

Problem

Based on the above considerations, an investigation was undertaken to establish which stretch training methods were superior in attaining the greatest range of active flexibility, within the session and over training weeks, with the most

enduring effects, and across different types of joints.

The purpose of this study was to examine active flexibility for:

A) Treatments

1) 3S (Scientific Stretching for Sports),
2) 2S (Static Stretching),
3) SS (Slow Stretching),
4) ES (Eccentric Stretching),
5) BS (Ballistic Stretching),
6) CS (Contract Relax Stretching),
7) Control (receiving no flexibility training);

B) Session

1) before session training,
2) after session training; and

C) Weeks

1) pre-training (pre-test),
2) two-week training,
3) four-week training,
4) six-week training (post-test),
5) two-week retention,

on shoulder extension and ankle plantar flexion.

Limitations of the Study

The study had the following limitations:

1) Extra-curricular activities were not controlled.
2) Subjects were restricted to right hand dominant, junior high school girls.

3) Subjects with joint impairment were excluded from the study.
4) Subjects received two different treatment conditions, one on the shoulder and one on the ankle.
5) The ankle joint (hinge) and shoulder joint (ball-and-socket) may not account for all the range of motion for ankle plantar flexion and shoulder extension.

Delimitations of the Study

The study had the following delimitations:

1) Subjects were confined to 42 female junior high physical education students, ranging from 12 to 14 years of age.
2) Joints selected, the right shoulder (ball and socket) and the left ankle (hinge), were of the synovial classification.

Definitions of the Study

Scientific Stretch for Sport (3S). This is a method of increasing flexibility whereby an isometric contraction of the muscles to be stretched (muscles in a lengthened position to start) is followed by a concentric contraction of the opposite muscle group, together with light force from a partner (external force), performed in a progressively more strenuous series until the range of movement cannot be increased without severe pain (Holt, 1973, pp. 5-7).

Static Stretch (2S). This is a method of increasing flexibility whereby an external force (i.e., hands pulling the segment) is applied in a progressively more forceful manner, until the range of movement cannot be increased, then held for a period of time, without severe pain (de Vries, 1976, pp. 436-437).

Slow Active Stretch (SS). This is a method of increasing flexibility by slow active concentric contraction of the agonist muscles, without additional assistance from an external force (i.e., hands pulling the segment to a greater range of motion), while relaxing the antagonist muscle group (the muscles to be stretched) (Jacobs, 1976, pp. 151-152).

Eccentric Stretch (ES). This is a method of increasing flexibility whereby a slow active eccentric contraction of the muscles to be stretched (muscles in a contracted position to start) is followed by light force from a partner (external force), while the subject relaxes both agonist and antagonist muscles around the joint.

Ballistic Stretch (BS). This is a method of increasing flexibility whereby particular emphasis is placed on training each individual to make all movements in a relaxed rhythmic manner, and to put special effort toward carrying the movement to the limit of the subject's range of motion (McCue, 1952, p. 322).

Contract Relax Stretch (CS). This is a method of increasing flexibility whereby an isometric contraction of the muscles to be stretched (muscle in a lengthened position to start) is followed by a concentric contraction of the opposite

muscle group (Moore and Hutton, 1980, p. 323).

Joint. This is a connection between two or more separate segments or parts of the skeleton. Joints are variously classified, with the official classification based on the material which joins the long parts. Classification of joints are: fibrous joint, cartilaginous joint, and synovial joint, with the latter being chosen for this study. This type of joint, which permits a variable amount of movement, consists of a series of investing ligaments, a true joint cavity, and hyaline cartilage over the articulating surfaces of the bones forming the joint. Synovial joints may be subdivided according to the type of movement: gliding, pivot, hinge, biaxial, and ball-and-socket joints (Francis, 1968). For the purposes of this study two sub-types of joints were chosen; shoulder (ball-and-socket) and ankle (hinge).

Shoulder Joint. This is the proximal ball-and-socket joint of the upper limb which controls the movement of the upper arm (humerus). Three basic movements of the shoulder are:

1) flexion and extension,
2) adduction and abduction, and
3) rotation.

Circumduction is merely a rhythmical combination in orderly sequence of flexion, abduction, extension, and adduction (Last, 1978). The present study termed any change in flexibility in shoulder extension, measured by the modified Leighton Flexometer, as occurring in the shoulder joint.

Ankle Joint. This is the distal hinge joint of the lower limb, which controls the movement of the leg relative to the foot, occurring in a sagittal plane. The ankle joint is not the only active joint when ankle plantar flexion becomes extreme; an increase in range is provided by hollowing of the plantar arches (Adams, 1971). The present study termed any change in flexibility in ankle plantar flexion as measured by the modified Leighton Flexometer as occurring in the ankle joint.

Shoulder Extension. This is the movement of the humerus posteriorly, in a sagittal plane (Last, 1978).

Ankle Plantar Flexion. This is the movement of the dorsum of the foot away from the anterior surface of the leg so that the foot tends to fall into line with the leg. This movement occurs through the sagittal plane (Adams, 1971).

CHAPTER II

REVIEW OF LITERATURE

Flexibility is the degree to which a joint(s) is free to move through its range of motion.

Overview of Flexibility

A review of the literature on flexibility revealed that there were several aspects of flexibility, which included: specificity, age, sex, strength, anthropometry, tests, limitations, somatotypes, activity patterns, warmup, motor performance, injury prevention and muscular soreness. The important area pertaining to the present study, flexibility training, will be discussed in greater detail (Song, 1979; Song and Reid, 1979; and Corbin and Noble, 1980).

Flexibility Training

Flexibility has been studied in connection with the science of sport, physical fitness and physical therapy. The following paragraphs review the literature on the effects of different flexibility training methods.

Moore and Hutton (1980) studied the effects of three different stretch techniques on passive hip flexibility for 23 female gymnasts, 17 to 23 years of age. Static stretch (S) required the subject to passively relax the hip extensors,

while a pulley and weight system was applied on the lower extremity in order to stretch the hip extensors. Contract-relax stretch (CR), a modified proprioceptive neuromuscular facilitation (PNF) technique, consisted of a five second maximal isometric hamstring contraction followed immediately by hamstring relaxation; the hip extensors were then stretched by the wall pulley apparatus. The contract-relax with agonist (hip flexors) contraction (CRAC) was identical to the CR procedures, with the adduction of a submaximal voluntary contraction of the agonists (hip flexors), while the pulley apparatus stretched the hamstring muscle group. The hip angles were measured with a goniometer and electrogoniometer. During one session, each subject performed three stretch techniques with three consecutive trials per stretch method. The performance order was a randomized latin square design. Data showed CRAC and S were equally superior in improving passive hip flexion over CR, during a one session test.

Medeiros, Smidt, Burmeister, and Soderberg (1977) studied the effects of different training methods on hip flexibility for 30 normal men ranging in age from 21 to 34 years. Subjects were randomly assigned to a control group (n = 10), passive stretch group (n = 10) and an isometric contraction group (n = 10). The training program consisted of eight treatment days. The passive stretch treatment group performed 20 passive stretches each treatment day at their threshold force, with the subject's right leg a total of six degrees beyond the threshold point. This position was maintained for three seconds before the subject's leg was returned to

the point where the threshold force was reached. Fifteen seconds of rest was allowed between each passive stretch. The isometric treatment group performed 20 isometric hip extension contractions each treatment day at the limb position where their threshold force was attained. The isometric contraction was held for six seconds with 15 seconds rest between each contraction. The control group received no exercise treatment. Before the first and after the last contraction or stretch on each treatment day the investigators measured the subject's pelvicfemoral angle at the point where the threshold force was attained. Stabilization was applied to the pelvis and lower left extremity. The data showed both treatments significantly increased the range of passive hip flexion compared to the control group. However, comparisons between the two treatment groups indicated that isometric contraction and passive stretch had similar effects.

Turner (1977) examined the effects of two different flexibility training methods, 3S (Scientific Stretching for Sport) and SS (Slow Active Stretching). 3S is a method of increasing flexibility whereby an isometric contraction of the muscles to be stretched (muscles in a lengthened position to start) is followed by a concentric contraction of the opposite muscle group, together with light pressure from a partner (external force) performed in a progressively more strenuous series until the scope of movement cannot be increased without severe pain. SS is a method of increasing flexibility by slow active concentric contraction of the agonist muscles without additional assistance from an external

force (i.e., hands pulling the segments to a greater range of motion), while relaxing the antagonist muscle group (the muscles to be stretched). Warm-up effects, different joint responses, and the standardization of training procedures were controlled. Subjects were 12 schoolgirl basketball players, aged 12 to 14. The research design consited of four replications of a 3 x 3 Graeco-latin square. Subjects were assessed for flexibility at the beginning of a six week flexibility training program, at the conclusion of training, and then after two two-week retention periods. Significant results showed that:

1) flexibility trained methods (3S and SS) improved flexibility.
2) the shoulder joint acquired more flexibility than either the knee joint of the ankle joint.
3) neither flexibility training method was superior to the other.
4) within two weeks of training cessation, both 3S and SS effects were lost to a significant degree when compared to control effects.
5) after four weeks of training cessation, there was no difference between changes of the control and two training groups.
6) there was no difference in loss of flexibility between the shoulder, the knee, and the ankle joints.

Song and Garvie (1976) examined 14 international Canadian freestyle wrestlers, ranging from 17 to 28 years of age. The study consisted of an experimental group (n = 8) and a control

group (n = 6). The flexibility program required the subjects to train for five weeks, four sessions per week. The 3S method of training involved the muscle group to be passively extended close to the maximum, then this same muscle group was isometrically contracted, against a fixed resistance, while in the extended position, for 5 to 7 seconds. The muscle group was then relaxed, thus making possible further extension of that particular group. With the antagonists relaxed while extended, further joint extension was allowed. Three exercises were performed on the shoulder, eight on the hip, one on the leg, one on the torso, and one on the combined trunk and hip. This program was undertaken in addition to the athletes' regular workout. Pre-post flexibility tests were taken on: trunk flexion, trunk lateral flexion, hip abduction, shoulder extension, hip extension, trunk extension, hip hyperextension, split pike, hip inward rotation, hip outward rotation, shoulder horizontal abduction and trunk rotation (19 tests). The experimental group showed all test results, except shoulder horizontal abduction, improved significantly. The control group showed 11 of the 19 test results improved significantly. However, between groups, only the left hip extension and split pike were significantly different.

Hartley (1976) investigated the effects of six methods of stretch on the active range of hip flexion for 119 women physical education students, ranging in age from 17 to 31 years. The program ran over a four week period with a total of 11 sessions. All subjects were tested before and after the study as well as before and after exercise on the first

day of each week. The treatment conditions were: passive lift and active hold, active PNF, ballistic and hold, relaxation method, passive PNF, and prolonged stretch. Passive lift and active hold required the subject's right leg to be lifted by the partner and taken slowly to the end-point (six seconds). The subject, at the end-point, held this position for the next six seconds by actively contracting the hip flexors. Passive pressure and active holding alternated every six seconds for one minute (five times). Active PNF required the subject to voluntarily lift the right leg as far into flexion as possible in six seconds. The subject, in the next six seconds, eased into a maximum isometric contraction down against the manual resistance of the partner. The subject alternated voluntary lift and maximum isometric extensor contractions so that each exercise occurred five times in one minute. Ballistic and hold required the subject to start at a 45 degree angle with four leg bounces made to flex the hip. On the fourth bounce an active hold was made to stop the leg at the end-point. Each leg bounce took one second, and the "hold" was maintained for six seconds. The whole combination was repeated five times in one minute. The relaxation method required the subject to use prolonged stretching with mental relaxation and mind-set techniques. Passive PNF required the right leg to be taken passively to the end-point in six seconds. Over the next six seconds, the subject eased into a maximum isometric contraction of the extensors against the maximal resistance of the partner. The passive lift and extensor contraction alternated so that each

was performed five times in one minute. The prolonged stretch required the right leg to be lifted passively and taken slowly to the end-point. The end position was held passively for one minute just below the pain threshold. The subjects were tested on the Leighton Flexometer for right hip flexibility. The data showed all methods increased flexibility significantly, but no one method was superior.

Tanigawa (1972) studied the effects of PNF hold-relax procedures, and passive mobilization on tight hamstring muscles for 30 normal subjects, ranging from 20 to 48 years of age. Subjects were randomly selected and assigned to a control group, a passive mobilization group, and a hold-relax group. Subjects were selected based on passive straight-leg-raising of less than 70 degrees in the right limb. The test measurement was a mathematical method involving the calculation of the sine of a right-angle triangle. Subjects were seen two days per week for four consecutive weeks with days one to six for attaining flexibility and days six to eight (one week) for retention of flexibility. All subjects were measured before and after treatment sessions. The passive mobilization technique required the stretching force (i.e., normally by the therapist) to be applied independently of any active movement by the subject. The hold-relax technique was used to increase joint range of motion and was based on an isometric contraction of the shortened muscle performed against maximal resistance. The results showed subjects receiving mobilization increased passive flexibility more than control subjects. The hold-relax procedure was superior to the passive mobilization

for developing and retaining flexibility. In addition, the hold-relax method increased flexibility at a faster rate than the passive mobilization.

Holt, Travis and Okita (1970) examined the effects of training on the hip extensor and lower back flexibility of 24 male college students. The investigation compared three methods - fast stretch (ballistic), slow stretch, and a modified version of PNF (IA-CA). This method required isometric contraction (5-6 seconds) of the agonist IA, followed by a concentric contraction of the antagonists, CA. Instead of diagonal patterns a simpler, unidimensional approach (movement and exercise in one plane) was adopted. This technique was based on the principles of successive induction, muscle relaxation, and reciprocal innervation. Six groups of four subjects each were formed and each group received all the treatments in different orders to control for treatment order effects. The program consisted of three weeks, three sessions per week. All methods were performed sitting and standing with a 30 second rest interval between positions, with the exception of IA-CA which was performed lying down instead of sitting. Each exercise was performed for 20 seconds, followed by rest for 10 seconds, for a duration of two minutes, with only one technique performed per week. The trunk flexion flexibility test was administered before and after each training session. The test designed by the author measured the distance subjects would reach forward while in a long sitting position. The data revealed the IA-CA method for increasing flexibility yielded significantly higher improvement scores

than both fast and slow stretch methods.

Fieldman (1966) examined the effects of selected extensibility exercises on the flexibility of the hip joint on 33 college students. The subjects were tested once a week for five weeks, with the exception of the first week, when tests one and two were performed. Each person was permitted one trial per test with the second test performed immediately after the first, the first serving as a warm-up. The third test was performed after the completion of four exercises and the fourth test was performed after the completion of six exercises. The fifth test followed the completion of eight exercises and the sixth test was performed within pre-exercise. All subjects were tested on a toe-touch test. The analysis showed all differences between tests were significant, with the exception of tests two and six, which showed that exercise improved the range of motion. The greatest gain occurred between tests two and three. The smallest gain was shown between tests four and five, suggesting subjects were reaching their maximum flexibility, but these tests still indicated greater total gain.

de Vries (1962) investigated the effects of seven 30-minute training sessions, over a five week period, with two training sessions per week, on the flexibility of 57 male college students. Subjects were randomly assigned to static and ballistic training groups. Static stretch involved a held position with no movement, slow or fast, in which the body segments to be stretched were locked into a position of greatest length. Ballistic stretch involved quick movements

characterized by quick jerks and pulls upon the body segments to be stretched. Static exercises (derived from Hatha Yoga exercises) started with a 30-second period and increased to one minute by the fourth workout. Ballistic exercises (derived from swim exercises) and static exercises started with 20 repetitions and increased to 40 repetitions by the fourth workout. Each group performed a total of eight exercises. Cureton's flexibility tests were employed: trunk flexion, trunk extension, and shoulder elevation. The pre-post test gains in flexibility for static and ballistic stretch methods showed trunk flexion, trunk extension, and shoulder elevation increased. The difference between static and ballistic flexibility gains revealed no significant difference.

Logan and Egstrom (1961) examined the relative effectiveness of slow and fast stretching techniques on the sacro-femoral angle for 12 female and 13 male university physical education students. Each sex group was randomly sub-divided into slow and fast groups. The slow-stretch method used the active contraction of the anterior muscle groups bent at the hips with the legs straight, maximally, and held the position momentarily. The fast-stretch method used momentum developed by a forced bobbing movement in which the weight of the trunk was bounced, against the resistance encountered at the range of motion. The test procedure required lines corresponding to the sacral indication and those formed by marks on the femur were then drawn on the photographs and the angle was measured with a protractor. The angle created reflected the

flexibility score. All subjects performed 20 repetitions per day. The subjects were measured again at the conclusion of the ten day exercise program. The results showed both slow and fast-stretch sub-groups had significantly increased the range of motion at the sacral-femoral angle. The females had increases significant at the 1% level for the fast-stretch sub-group and at the 5% level for the slow-stretch sub-group. The males had increased significantly at the 1% level for both sub-groups. There was no significant difference between the mean difference of the final measurement of the fast and slow-stretch for either the males or the females. The conclusion drawn was that there was no difference between slow and fast-stretching.

McCue (1952) determined the effects of three weeks of stretch exercises on the flexibility for the lower quartile of 130 college women, ranging in age from 17 to 26 years (mean age 19.5 years). The bounce technique was performed in a relaxed rhythmic manner, to the limit of range of motion. The exercises and tests involved the following joints: hip flexion, hip rotation, ankle flexion and extension, trunk flexion and total back and neck extension. The measurements were taken with the goniometer just before the exercise program, immediately following the three-week series, and again eight weeks following the series. All mean differences between all pre-post test scores on the three-week training period were significant. In addition, hip flexion, total back and neck extension, and hip rotation retained significant amounts of flexibility eight weeks following the

exercise program.

Weber and Kraus (1949) investigated the effects of the spring stretch method (bobbing) on group one (n = 25 cases) and plain stretch method (control) on group two (n = 25 cases) for the toe-touch and leg-lift from supine position tests and exercises. The passive spring stretch of hamstring muscles was described as a "bobbing" motion caused by the physical therapist at the point where it is felt that the muscle has reached its fullest length. The active spring stretch is done with the patient standing erect, knees straight, bending over to touch the floor with fingertips. The subject bobs up and down with a spring-like motion, exerting power in the downward direction, thus decreasing the distance from fingertips to the floor. The plain stretch applied was the same exercise, but without the bobbing or spring-like rhythmical motion. The maximum range of motion was held for a few seconds. The rate of improvement per month in regaining normal length of these muscle groups was computed in 50 unselected cases. The technique of measurements was both degree (protractor) and linear (inches) for the leg-lift and toe-touch respectively. The authors reported the spring stretch method showed an average improvement of 3.0 degrees in the hamstring muscles (leg-lifts) and 2.0 inches in the back-hamstring-gastrosoleus muscles (toe-touch) per month. The plain stretch improved by an average of 1.0 degree and 1.0 inch per month respectively. The final conclusion was that spring stretch method showed 200 percent more efficiency in the hamstring muscle (leg-lift) stretch and 100 percent more efficiency in

the back-hamstring-gastrosoleus muscle (toe-touch) stretch.

Summary

Table 1 shows a summary of the results of different studies conducted on flexibility training.

The major generalizations from these studies are:

1) Flexibility training, regardless of method, has increased active and passive flexibility within one training session and over weeks of training.
2) Active and passive flexibility retention have been retained for at least one week following cessation of flexibility training.

The generalizations across different flexibility training studies are few, due to the following limitations:

1) Researchers employing the same stretch technique have not followed standard procedures.
2) Test methods utilized by different investigators (i.e., linear and degree) may not be comparable, since linear change may be shown without a change in degrees.
3) Immediate training effects and long term training effects are different; generalizations made should be specific to each effect. Research papers examining both effects have failed to report the results in this manner.
4) Active flexibility and passive flexibility are distinctly different; information about one type should not be transferred to the other. Some research papers investigating different joint flexibility measurements have not

TABLE I

EFFECTS OF DIFFERENT STRETCH TRAINING METHODS ON FLEXIBILITY

Author	Subjects	Sex	Weeks	Sessions	Method	Flexibility Type	Test B	Test A	Flexibility Test	Results: Training	Results: Retention
Moore et al. 1958	33	F		1	CR CHAC S	Passive	No	Yes	Goniometer (°)	Improvement CHAC and S superior	No data
Pedeiros 1977	36	M		6	Passive Isometric contraction	Passive	Yes	Yes	Goniometer (°)	Improvement No difference bet. methods	No data
Turner 1977	12	F	10	30	SS SS	Active	No	Yes	Leighton Flexometer (°)	Improvement No difference	Retention No difference
Hartley 1976	110	F	6	11	Passive lift and active hold Ballistic hold Active PNF Relaxation Passive PNF Prolong stretch	Active	Yes	Yes	Leighton Flexometer (°)	Improvement No difference bet. methods	No data
Gang et al. 1976	15	M	6	30	SS	Active (?)	No	Yes	Leighton Flexometer (°) (?)	Improvement Left hip extension and split pike superior to control	No data
Yoshimo 1972	30	M	6	6	Hold-relax Passive	Passive	Yes	Yes	Path method (°)	Improvement Hold-relax superior	Retention Hold-relax superior
Holt et al. 1970	24	M	3	9	Ballistic Slow IA-CA	Active	Yes	Yes	Sit and Reach (in)	Improvement IA-CA superior	No data
Fieldman 1966	33	M	5	6	Exercises	Active	Yes	Yes	Toe touch Test (in)	Improvement	No data
de Vries 1962	57	M	8	7	Static Ballistic	Active	No	Yes	Curton's method (in)	Improvement No difference bet. methods	No data
Lepes et al. 1961	35	F/M	1 1/2	18	Slow Fast	Active	No	Yes	Goniometer (°)	Improvement No difference	No data
McCue 1952	110	F	3	9	Bounce	Active	No	Yes	Goniometer (°)	Improvement	Retention
Weber et al. 1949	18	M/F	8		Spring Plain	Passive Active	No	Yes	Goniometer (°) Toe-touch Test (in)	Improvement Spring superior	No data

Test B: before and after test within the training session
Test A: tests over weeks
(?): incomplete
(°): degrees

clearly indicated which type of flexibility was studied.

5) Studies examining a treatment effect without the use of a control group seriously affect the interpretation of the results.

6) Training programs without proper supervision may not guarantee the performance of the program.

7) Joint tests with significantly higher reliability than other joint tests have greater power to detect treatment changes; comparison between these joint tests may not be justified.

8) Since joints are unique within the across-joint-classification, the response to treatment may be different. Consequently, pooling results across joints may prove meaningless.

9) Research has yet to examine a number of standard stretch techniques in one study. An attempt has been made to investigate a number of combined methods (i.e., ballistic - hold versus passive lift - active hold). The superiority of one method cannot be claimed.

10) Since men and women have both physiological and anatomical differences within joints (i.e., hips), their response to treatment may not be the same.

11) The presence or absence of a pre-test warm-up in different studies makes the results among studies difficult to compare (Atha-Wheatley, 1976).

12) Little attention has been given to equalize time in training from one method to the next. Perhaps one method may prove more or less effective only because it requires more

or less time in training.

13) The subject's initial flexibility score can affect the rate and magnitude of flexibility change. Researchers have neglected to allow for this important variable. If the study had blocked on initial flexibility scores, their results were not reported in this manner.

14) The frequency and duration of training sessions over days and/or weeks are extremely different from one study to the next. Training programs have little in common for cross comparisons.

It may therefore be concluded that previous investigations were not able to discern which flexibility training method(s) was superior in developing both immediate and long term flexibility across different joints, with the most enduring effects. Consequently, a major comprehensive investigation on flexibility training was designed and undertaken with a view to minimize the effects of the limitations cited above.

CHAPTER III

METHODOLOGY

Subjects

Forty-nine volunteer female, right hand dominant, junior high students, ranging from 12 to 14 years of age, at Lakewood Intermediate School, Kenora, Ontario were selected as subjects. Permission to test the subjects was received from the principal of the school (Appendix C). The preferred writing hand was used to categorize the subjects as right-handed.

Forty-nine shoulder joints and 49 ankle joints were studied, each subject contributing both joints. Subjects with any history of joint injury or disease (e.g., broken ankle) were excluded from the study.

Randomization

Selection. Two classes totalling forty-nine female subjects were selected from a school population of 350 females.

Assignment. Following the initial flexibility tests, each subject was rank-ordered on her joint flexibility scores, from high to low, for each joint. Blocks of seven subjects were formed by ranking successive students on flexibility. Subjects in each block were then randomly assigned to one of the seven conditions. Assignments to the six treatment and

control conditions were done independently for ankle and shoulder joints except that in the case of the control conditions, no subject was allowed to be in the control group for both the shoulder and ankle studies.

Modification. Three subjects dropped out of the study, leaving six subjects in each of three treatment groups and seven subjects in each of the remaining four treatment groups. Since this occurred, one subject's joint was randomly selected from each of the remaining four groups and discarded from the analysis. This procedure was performed independently for both the ankle joint and the shoulder joint. Subsequently, a one-way analysis of variance among treatments was conducted on pre-test flexibility scores (Cook and Campbell, 1979, p. 152). The analysis was performed independently for both the ankle and shoulder joints. The results showed no significant difference among treatment groups for the pre-test flexibility scores for each ankle and shoulder joint analysis ($p > 0.05$).

Experimental Design

The experimental design consisted of two 7 x 5 x 2 (treatment, week and session) factorial designs with repeated measures on the last two factors.

The first factor (factor A) was the treatment variable. The seven levels of this factor were:

A_1 3S training

	Weeks:	B_1		B_2		B_3		B_4		B_5	
	Subjects	Sessions: C_1	C_2	C_1	C_2	C_1	C_2	C_1	C_2	C_1	C_2
A_1	1 2 6										
A_2	1 2 6										
A_3	1 2 6										
A_4	1 2 6										
A_5	1 2 6										
A_6	1 2 6										
A_7	1 2 6										

Treatment Conditions

A_1 3S training (n = 6)
A_2 2S training (n = 6)
A_3 5S training (n = 6)
A_4 8S training (n = 6)
A_5 8S training (n = 6)
A_6 CS training (n = 6)
A_7 Control (n = 6)

Week in Experiment

B_1 pre-training
B_2 two-week training
B_3 four-week training
B_4 six-week training
B_5 two-week retention

Session

C_1 pre-session
C_2 post-session

Figure 3. Design of the study.

A_2 2S training

A_3 SS training

A_4 ES training

A_5 BS training

A_6 CS training

A_7 control

The second factor (factor B) involved repeated measures. The five levels of this factor were:

B_1 pre-training (pre-test)

B_2 two-week training

B_3 four-week training

B_4 six-week training (post-test)

B_5 two-week retention

The third factor (factor C) also involved repeated measures. The two levels of this factor were:

C_1 pre-session training

C_2 post-session training

Testing Apparatus

The Leighton Flexometer was developed for measuring the shoulder joint and ankle joint for active flexibility. The flexometer is comprised of a weighted 360 degree dial and a weighted pointer housed in a case. The dial and pointer function freely and independently of each other, though each is controlled by gravity. The flexometer, when positioned within 20 degrees off the horizontal, will record accurate angular movement. Independent locking mechanisms are furnished

for the dial and the pointer which stop angular movement of either at any position. The flexometer must be strapped to the segment next to the joint being tested. The dial will be locked at one position (e.g., full flexion of the ankle); the angular movement will be performed and the pointer locked at the opposite extreme position (e.g., full extension of the ankle). The exact reading of the pointer on the dial will be the angular movement through which motion will have taken place (Leighton, 1966).

The Leighton Flexometer test demonstrated the correlation coefficient between the first and second measurement to be 0.913 and 0.996 (thirty measures on 120 boys) (Leighton, 1966, p. 86). Joint tests with significantly higher reliability than other joint tests have greater power to detect treatment changes; comparison between these joint tests may not be justified. Consequently, the author developed the modified Leighton Flexometer test method for ankle plantar flexion and shoulder extension to alleviate this problem.

The reliability of two different flexibility test methods, the traditional Leighton Flexometer method and a modified Leighton Flexometer method was investigated. Twenty-four female volunteer physical education students, ranging in age from 18 to 25 years, were selected as subjects. Twelve of these subjects were assigned to the traditional flexometer method and 12 to the modified Leighton Flexometer testing method. Following the orientation session, in which subjects were familiarized with the testing technique, three tests were conducted during the second session. The results of the

last two tests were analyzed statistically using the Pearson product moment correlation coefficient and the difference between two correlation coefficients for independent samples was tested.

Analysis of data revealed that the traditional Leighton Flexometer method gave correlation coefficient values of .914 for shoulder extension ($p < 0.01$) and .990 for ankle plantar flexion ($p < 0.01$). The modified Leighton Flexometer method on the other hand revealed correlation values of .992 and .993 respectively ($p < 0.01$). Differences between the methods for the correlation coefficients for the shoulder joint were significant ($p < 0.01$) and for the ankle joint the difference was not significant ($p > 0.05$). The modified Leighton Flexcmeter was used in this study because it showed a reliability of .99 across both joints.

Modified Leighton Flexometer. The original cloth strap was removed from the Flexometer. A lightweight (28.35 g) preshaped, plastic brace (foam lined) with adjustable elastic-velcro strap (195.5 to 241.3 mm) was attached to the back of the instrument. This modification allowed for the instrument to fit any size arm or foot comfortably (Figures 4 and 5).

The modified Leighton Flexometer method was consistently more reliable for shoulder extension and ankle plantar flexion tests than the traditional Leighton Flexometer method. This has allowed for better comparison across joints, when detecting



Figure 4. Front view of the modified Leighton Flexometer.



Figure 5. Back view of the modified Leighton Flexometer.

treatment changes.

Apparatus. All testing was carried out in a centrally heated gymnasium. The testing room was at a temperature between 18 and 22°C on all testing days. The room was equipped with two portable tables for testing the shoulder (Figure 6) and ankle joints (Figure 7). One adjustable strap and one cylinder were used for securing subjects into the proper ankle joint test position (Appendix B). Subjects wore standard physical education clothing (e.g., a pair of shorts, a T-shirt, and bare feet).

Testing Procedures

Leighton's rules for flexibility testing (Leighton, 1966) were followed, with modification, as a standard procedure for each test. All tests were conducted in the sagittal plane, in one direction per joint: shoulder extension and ankle plantar flexion. The flexometer's dial was permanently locked, by horizontally positioning the instrument on the table, since only one direction of flexibility was to be measured.

The initial flexibility testing was undertaken within one week before the first flexibility training session. A post-training test was given during the last training session. The retention test occurred two weeks after the post-training test. The flexibility training tests were held on regular school days between 8:30 a.m. and 4:30 p.m.



Figure 6. Shoulder extension table.



Figure 7. Ankle plantar flexion table.

The pre-training, training, post-training and retention tests all followed the same procedures. In each test session two subjects were randomly called for testing. They changed into their physical education clothing and moved directly to the testing room. Upon completion of the shoulder and ankle tests, the subjects left the testing room and two more subjects were called. This pattern was continued until all subjects had been tested.

The investigator collected all the data for this study. The tester was helped by an assistant who recorded the subjects' flexibility scores for each joint being tested. All subjects performed three consecutive maximum movements for each specific joint to be tested, as instructed by the tester. Each test trial was three seconds in duration. Each subject was given fifteen seconds rest between test trials.

The flexometer was checked before each test session to ensure it was in working order.

Flexibility Tests

The shoulder extension test (Figure 8) and ankle plantar flexion test (Figure 9) were administered to each subject. For a further explanation on each test see Appendix B.

Training Apparatus

All training was carried out in a large centrally heated school gymnasium. The temperature of the training gymnasium



Figure 8. Shoulder extension test.



Figure 9. Ankle plantar flexion test.

was between 18°C and 22°C during all training sessions. Subjects wore standard physical education clothing (e.g., a pair of shorts, a T-shirt, and stockings, or bare feet).

Training Procedures

The specific joint training method was followed as a standard procedure. Subjects were encouraged to perform only the joint method assigned. During the study, the subjects were asked to refrain from any flexibility training other than that associated with the experiment. A reward system was implemented to provide motivation for subjects to meet this instruction. A movie and lunch (Big Mac, apple and milk) was provided for all students and staff members involved after the fourth week of the training study. In addition a plaque from the University of Alberta was presented to the principal of Lakewood Intermediate School in appreciation for his school's full co-operation during the course of the experimental research.

The flexibility training program consisted of 24 training sessions. Each exercise was performed three consecutive times during the session. Each subject chose one permanent partner to assist when necessary in the exercises. The order of the joints to be exercised was changed in each series.

The workload for each subject was standardized:

1) thirty seconds training per session(s), and
2) fifteen seconds rest intervals.

Each subject received two treatment conditions:

1) on the shoulder joint, and

2) on the ankle joint.

The exercises were performed during each physical education class. The gymnasium was divided into six stations: 3S, 2S, SS, ES, BS, and CS. The distance between the stations was great enough so that subjects exercising at each station did not distract one another. During each training session, the instructor supervised the training to further ensure that the exercise was performed properly. The instructor accomplished this by moving from station to station, observing, assisting, and encouraging the subjects.

Flexibility Training

The present study developed standard procedures for the six different flexibility training methods (3S, 2S, SS, ES, BS, and CS) for shoulder extension and ankle plantar flexion. The following is an explanation on each standardized flexibility training method. For further explanation see Appendix A.

Scientific stretch for sport training (3S). A maximal range of motion is attained by actively and passively flexing or extending a segment nearest to the joint being trained, in an arc, as far as possible. Repetitions of the exercise are performed, without severe pain. One repetition consists of:

a) two second concentric contraction,

b) six second isometric contraction, and

c) two second concentric contraction, with light

force from a partner.

Subject repeats the above exercise (rest optional). The contractions performed by the exercise subject must be gradually, not explosively, increased in effort.

Static stretch training (2S). A maximal range of motion is attained by passively flexing or extending a segment nearest to the joint being trained, in an arc, as far as possible. Repetitions of the exercise are performed without severe pain. One repetition consists of:

a) two second passive extension, and

b) eight second held passive extension.

Subject relaxes for fifteen seconds before repeating each exercise. The external force applied to extend the segment is gradually, not explosively, increased in effort.

Slow active stretch training (SS). A maximal range of motion is attained by actively flexing or extending a segment nearest to the joint being trained, in an arc, as far as possible. Repetitions of the exercise are performed, without severe pain. One repetition consists of:

a) two second concentric contraction, and

b) eight second isometric contraction.

Subject relaxes for fifteen seconds before repeating each exercise. The contractions performed by the subject must be gradually, not explosively, increased in effort.

Eccentric stretch training (ES). A maximal range of motion is attained by actively flexing or extending a segment

nearest to the joint being trained, in an arc, as far as possible. Repetitions of the exercise are performed, without severe pain. One repetition consists of:

a) five second eccentric contraction, and

b) five second passive extension.

Subject relaxes for fifteen seconds before repeating each exercise. The contraction and passive extension performed by the subject must be gradually, not explosively, increased in effort.

Ballistic stretch training (BS). A maximal range of motion is attained by actively flexing or extending, in a relaxed rhythmic manner, a segment nearest to the joint being trained, in an arc, as far as possible. Repetitions of the exercise are performed, without severe pain. One repetition consists of one second concentric contractions. Subjects will perform 10 consecutive repetitions, then relax for fifteen seconds before repeating each exercise. The series of ten concentric contractions is performed in a rhythmic manner, explosive in effort.

Contract relax stretch training (CS). A maximal range of motion is attained by actively flexing or extending a segment nearest to the joint being trained, in an arc, as far as possible. Repetitions of the exercise are performed, without severe pain. One repetition consists of:

a) two second concentric contraction,

b) six second isometric contraction, and

c) two second concentric contraction.

Subject immediately repeats the above exercise (rest optional). The isometric contraction performed by the exercise subject must be gradually, not explosively, increased in effort.

All subjects who performed flexibility training on the shoulder joint were required to flex at the knee about 90 degrees (Figure 10) and any partner who assisted a subject was required to stay on her feet or buttocks (Figure 11). This procedure was followed to avoid stretching of the ankle joint, thereby causing a training effect.

Summary of Test and Training Format

Pre-test orientation. The subjects were familiarized with both the training and test methodology. Test and training methods were presented on slides for subjects to follow standard procedures. An information sheet pertaining to shoulder or ankle injury was completed by the students.

Initial test. The initial flexibility test was one week before the first training session.

Training program. The flexibility training sessions were held three times per week (i.e., Monday, Wednesday, and Thursday) for six consecutive weeks, in conjunction with the physical education classes.

Training tests. Two tests were administered per joint, one before and one after the training session. The subjects



Figure 10. Subject is flexed at knee.



Figure 11. Partner is supported by feet.

were assessed for active flexibility at the first, sixth, twelfth, eighteenth and twenty-fourth session of the training program.

Post-training period. The subjects did not receive any flexibility training during the seventh and eighth week of the study. The physical education classes continued as usual.

Post-training test. The subjects were assessed after a two-week retention period (no flexibility training) (Figure 12).

Week	Session	Test
0		Pre Post T_0
1	1	T_1
	2	
	3	
2	4	
	5	
	6	T_2
3	7	
	8	
	9	
4	10	
	11	
	12	T_3
5	13	
	14	
	15	
6	16	
	17	
	18	T_4
7	19	
	20	
	21	
8	22	
	23	
	24	T_5
9		

Figure 12. Schematic diagram of testing and training format.

CHAPTER IV

RESULTS AND DISCUSSION

The ankle data and shoulder data were analyzed separately using a three-way analysis of variance with repeated measures on the last two factors (Winer, 1971, pp. 539-559). A computer program ANOV34 developed by the Division of Educational Research Services at the University of Alberta was used (DERS, 1979). For both analyses, the training factor (factor A) had seven levels:

A_1 3S training

A_2 2S training

A_3 SS training

A_4 ES training

A_5 BS training

A_6 CS training

A_7 Control

The first repeated measure (factor B) consisted of the five sessional measurements made over the training period.

B_1 pre-training (pre-test)

B_2 two-week training

B_3 four-week training

B_4 six-week training (post-test)

B_5 two-week retention

The second repeated measure (factor C) consisted of pre- and post-session measures taken at each of the measurement sessions in factor B.

C_1 pre-session training

C_2 post-session training

In testing effects involving repeated measures, the Greenhouse-Geisser procedure was used (Winer, 1971, p. 523). This procedure guards against potential violations of the homogeneity of covariance assumptions.

Ankle Plantar Flexion

The summary analysis of variance for ankle plantar flexion is shown in Table 2. The cell means are displayed graphically in Figure 13. Significant main effects were found for B (week) and C (within-session). A significant treatment by within-session-interaction was also noted. Application of the Greenhouse-Geisser conservative test yielded conclusions identical to those in the regular test. Neither the treatment main effect nor any other interaction was significant.

Scheffé comparisons were conducted on the ankle joint data. In the case of factor B (Table 3 and Figure 14), flexibility significantly increased between the fourth and sixth week and then decreased after the two week retention period. Subjects' mean range of motion after the retention period was not significantly different from the pre-test scores.

In the case of factor C, the average post-session mean was significantly higher than the average pre-session mean ($\bar{X}$ pre = 106.3 and $\bar{X}$ post = 107.9).

The differential treatment effect as revealed by the AC interaction showed that 3S, 2S, ES and CS significantly increased within-session flexibility when compared to the control (Table 4 and Figure 15). In addition, 3S training was superior to CS training in relation to within-session gains (Table 5).

Shoulder Extension

The summary of analysis of variance for shoulder extension is shown in Table 6 with cell means displayed in Figure 16. As in the ankle data only B (week), C (within-session) and AC (treatment by within-session) were significant. Application of the Greenhouse-Geisser test did not change the conclusions.

Scheffé comparisons were conducted on the shoulder joint data. In the case of factor B (Table 7 and Figure 17), flexibility significantly increased between the beginning and second week, between the fourth and sixth week and then decreased after the two week retention period. Subjects' mean range of motion after the retention period was significantly different from the pre-test scores.

In the case of factor C, the average post-session mean was significantly higher than the average pre-session mean ($\bar{X}$ pre = 161.6 and $\bar{X}$ post = 162.5).

The differential treatment effect as revealed by the AC interaction showed that only 3S significantly increased within-session flexibility when compared to control (Table 8 and

TABLE 2

SUMMARY OF ANALYSIS OF VARIANCE ON THE ANKLE DATA

Source of Variation	Sum of Squares	Degrees of Freedom	Mean Squares	F Ratio
Between Subjects	24869.00	41		
Factor A (treatment)	1023.00	6	170.50	0.25
Subject Within Group	23846.00	35	681.31	
Within Subjects	3159.00	378		
Factor B (week)	1237.00	4	309.25	41.59**
Factor A x Factor B (AB)	203.00	24	8.46	1.14
Factor B x Subjects Within Group	1041.00	140	7.44	
Factor C (session)	277.00	1	277.00	277.00**
Factor A x Factor C (AC)	232.00	6	38.67	38.67**
Factor C x Subjects Within Group	35.00	35	1.00	
Factor B x Factor C (BC)	8.00	4	2.00	2.35
Factor A x Factor B x Factor C (ABC)	7.00	24	0.29	0.34
Factor B x Subjects Within Group	119.00	140	0.85	

Greenhouse-Geisser Conservative Test

* Significant at 5% level ($p < 0.05$)
** Significant at 1% level ($p < 0.01$)



Figure 13. Mean flexibility scores for each treatment, week, and session on the ankle joint.

TABLE 3

SCHEFFÉ POST-HOC CONTRASTS AMONG LEVELS OF FACTOR B (WEEK) ON THE ANKLE DATA

Comparison	F Ratio
B_1, B_2 (Week 1, Week 2)	0.00
B_1, B_3	0.09
B_1, B_4	56.95*
B_1, B_5	2.77
B_2, B_3	0.10
B_2, B_4	57.20*
B_2, B_5	2.83
B_3, B_4	52.48*
B_3, B_5	1.85
B_4, B_5	34.61*

* Significant ($p < 0.05$)



Figure 14. Mean flexibility scores for each week on the ankle joint.

TABLE 4

SCHEFFÉ POST-HOC CONTRASTS AMONG LEVELS OF FACTOR C (SESSION) FOR GIVEN LEVELS OF FACTOR A (TREATMENT) ON THE ANKLE DATA

Treatment	Comparison	F Ratio
A_1 (3S training)	C_1, C_2 (pre- and post-session)	81.40*
A_2 (2S training)	C_1, C_2	53.40*
A_3 (SS training)	C_1, C_2	1.71
A_4 (ES training)	C_1, C_2	48.17*
A_5 (BS training)	C_1, C_2	2.04
A_6 (CS training)	C_1, C_2	17.00*
A_7 (control)	C_1, C_2	0.14

* Significant ($p < 0.05$)



Figure 15. Mean flexibility scores within-session for each treatment on the ankle joint.

TABLE 5

SCHEFFÉ POST-HOC CONTRASTS AMONG LEVELS OF FACTOR A (TREATMENT) FOR SIGNIFICANT C_1 AND C_2 DIFFERENCES ON THE ANKLE DATA

Comparison	F Ratio
A_1 (C_2-C_1), A_2 (C_2-C_1) (3S, 2S)	5.25
A_1 (C_2-C_1), A_4 (C_2-C_1) (3S, ES)	6.38
A_1 (C_2-C_1), A_6 (C_2-C_1) (3S, CS)	15.00*
A_2 (C_2-C_1), A_4 (C_2-C_1) (2S, ES)	1.13
A_2 (C_2-C_1), A_6 (C_2-C_1) (2S, CS)	9.75
A_4 (C_2-C_1), A_6 (C_2-C_1) (ES, CS)	8.63

* Significant ($p < 0.05$)

TABLE 6

SUMMARY OF ANALYSIS OF VARIANCE ON THE SHOULDER DATA

Source of Variation	Sum of Squares	Degrees of Freedom	Mean Squares	F Ratio
Between Subjects	26823.00	41		
Factor A (treatment)	5435.00	6	905.83	1.48
Subject Within Group	21388.00	35	611.09	
Within Subjects	33144.00	378		
Factor B (week)	21864.00	4	5466.00	82.71**
Factor A x Factor B (AB)	784.00	24	32.67	0.49
Factor B x Subjects Within Group	9252.00	140	66.09	
Factor C (session)	77.00	1	77.00	13.08**
Factor A x Factor C (AC)	204.00	6	34.00	5.78**
Factor C x Subjects Within Group	206.00	35	5.89	
Factor B x Factor C (BC)	42.00	4	10.50	2.15
Factor A x Factor B x Factor C (ABC)	58.00	24	2.42	0.49
Factor B x Subjects Within Group	685.00	140	4.89	

Greenhouse-Geisser Conservative Test

* Significant ($p < 0.05$)
** Significant ($p < 0.01$)



Figure 16. Mean flexibility scores for each treatment, week, and session on the shoulder joint.

TABLE 7

SCHEFFÉ POST-HOC CONTRASTS AMONG LEVELS OF FACTOR B (WEEK) ON THE SHOULDER DATA

Comparison	F Ratio
B_1, B_2 (Week 1, Week 2)	47.93*
B_1, B_3	69.25*
B_1, B_4	159.49*
B_1, B_5	56.74*
B_2, B_3	1.95
B_2, B_4	32.55*
B_2, B_5	0.37
B_3, B_4	18.55*
B_3, B_5	0.62
B_4, B_5	25.98*

* Significant ($p < 0.05$)



Figure 17. Mean flexibility scores for each week on the shoulder joint.

TABLE 8

SCHEFFE POST-HOC CONTRASTS AMONG LEVELS OF FACTOR C (SESSION) FOR GIVEN LEVELS OF FACTOR A (TREATMENT) ON THE SHOULDER DATA

Treatment	Comparison	F Ratio
A_1 (3S training)	C_1, C_2 (pre- and post-session)	13.46*
A_2 (2S training)	C_1, C_2	0.63
A_3 (SS training)	C_1, C_2	1.02
A_4 (ES training)	C_1, C_2	3.49
A_5 (BS training)	C_1, C_2	0.07
A_6 (CS training)	C_1, C_2	0.50
A_7 (control)	C_1, C_2	0.08

* Significant ($p < 0.05$)



Figure 18. Mean flexibility scores within-session for each treatment on the shoulder joint.

Figure 18).

Discussion of Treatment Over Time

Analysis of the data on both the ankle and shoulder joints revealed no differential effect of treatments over the weeks of training when compared to the control group. There can be several explanations for this. First, control subjects training on one joint may have had a learning effect on the control joint; that is, the subject, after having worked on the other joint may have realized that she could extend the control joint even further. Second, the flexibility test itself may have had an effect on the control subjects' joints, since the control subjects received nine seconds of stretching on both joints within each test session. Third, all subjects including control subjects received 40 minutes of physical education per day during which volleyball classes were introduced between the beginning and second weeks and gymnastics classes between the fourth and sixth weeks. These physical activities required subjects to stretch the shoulder joint during the volleyball sessions (i.e., underhand bump skill) and increased both shoulder and ankle joint flexibility during the gymnastics sessions (i.e., floor routines). This may have had an effect on flexibility. The results showed that all groups increased in shoulder extension during the first two weeks and increased both shoulder extension and ankle plantar flexion for the last two weeks of training. During the fourth and fifth weeks there was no effect. Fourth, some control members were

involved in an intramural program (e.g., volleyball), extra-mural program (e.g., gymnastics), or extra-curricular activities (e.g., competitive swimming). The nature of these activities again required subjects to stretch both ankle and shoulder joints maximally and this may have caused increased flexibility in the control group. Fifth, the amount of training each joint received may not have been of sufficient magnitude to cause major flexibility development. Each joint received 30 seconds of training per session, one-and-a-half minutes per week, three minutes every two-week test interval for a total of 12 minutes of training, over the entire six week training program. Perhaps both the ankle and shoulder joints needed sufficiently more time in flexibility training to allow for the mechanisms underlying change in range of motion to have a sustaining effect. Unfortunately, "there are no research studies which specify the minimal amount of time required for effective stretching" (Trombly and Scott, 1980, p. 226).

Past research revealed other studies which have shown that the control group increased joint range of motion over weeks of investigation. Tanigawa (1972) showed the control group increased in hip flexion. He attributed this result to the halo effect. Tanigawa explained this effect as being the result of passively extending the hip joint on each test occasion and that flexibility became a major concern of the control. Song and Garvie (1976) reported that the control group increased in flexibility on a number of tests, over four weeks of wrestling training. The control groups' gains in flexibility was attributed to regular stretching and wrestling

training. Hartley (1976) showed improved range of motion for the hip joint for the control group. She believed the conditioning activities that the subjects underwent during daily physical education and test sessions influenced the control's joint scores. Turner (1977) observed control increased in flexibility throughout the entire study. He suggested the nature of the sport activity and athletes' active lifestyles influenced the control subjects' flexibility scores.

From present and past research, the control group's range of motion could have easily been affected by a number of extraneous variables. In addition, the time in joint sessional training may not have been of sufficient duration to cause a differential effect of treatments over the weeks of training when compared to the control group. The author suggests that future research on this area make two major adjustments. First, use different durations of time for training with none less than 30 seconds. Second, subjects chosen should have very little physical activity (e.g., sedentary 25-30 year olds). Moreover, research studies showing changes in range of motion, without comparison to a control group, cast serious doubt as to the effectiveness of the methods (e.g., de Vries, 1962 and McCue, 1952).

Joint differences for within-session training. The analyses on the ankle and shoulder data for factor C (within-session training) were not identical. A comparison of the ankle (Figure 13) and the shoulder (Figure 16) showed the responses for within-session training at each repeated measure were

similar, with the exception of the fourth repeated measure on the shoulder joint. The graph (Figure 16) on the shoulder suggested that the subjects involved in the 2S, ES and CS methods reached a ceiling effect (175 degrees), while the 3S subjects did not encounter this problem. The ceiling effect may have accounted for some of the dissimilar results for the ankle and shoulder for within-session flexibility training.

Since the 3S method seemed to produce the greatest within-session flexibility increase in both the ankle and shoulder studies it is worthwhile to examine it a little more closely.

The following is a review of the scientific investigations pertaining to 3S training and a theoretical explanation on neural mechanisms underlying stretch training which may help explain the findings of the present study.

Scientific Investigations on 3S

Holt (1973) supported the effectiveness of increased range of motion for the 3S training method based on two research studies: Holt, Travis and Okita (1970) and Tanigawa (1972).

Holt, Travis and Okita (1970) studied the effects of three stretching techniques for increasing hip extension and low back flexibility. Fast-stretch (ballistic), slow-stretch, and IA-CA - a modified version of PNF. IA-CA is defined as an isometric contraction of the agonist (IA), followed by a concentric contraction of the antagonist (CA). Results showed

that the IA-CA stretch technique had maximum effect on flexibility.

Tanigawa (1972) examined the effects of PNF and passive mobilization on passive range of motion for tight hamstring muscles. The investigation revealed that the PNF method of therapeutic exercise resulted in a significant increase in flexibility along with rate of increase compared to passive mobilization.

A close examination of the above procedures revealed these methods were distinctively different from the 3S procedure defined by Holt (1973). Holt et al. (1970) described the IA-CA method as

> E [partner] when assisting, did not push or apply force to the leg in any direction, but served merely as a stationary object against which S could apply force for contraction of the hip extensors (p. 614).

The 3S training method states that the partner, after a six second isometric contraction is to assist the subject with light force in the direction the subject is concentrically contracting the opposite muscle group to be stretched.

Tanigawa (1972) described the PNF technique as the therapist passively lifting the patient's limb through multiple plane and joint movements. The therapist served as an immovable object so that the patient could perform isometric contractions, followed by only passive extension of the joint. This method is not in accordance with Holt's 3S procedures.

Based on the above considerations, the methods described by Holt et al. (1970) and Tanigawa (1972) are very different from the procedures defined as 3S training. The relative

effectiveness of IA-CA training and the PNF techniques respectively, cannot be generalized to the 3S training method.

Since the claims made by Holt (1973), additional research on 3S training methods has been undertaken (Song and Garvie, 1976 and Turner, 1977).

Song and Garvie (1976) compared the flexibility of wrestlers using the 3S flexibility method, with that of a control group receiving no 3S training. The experimental group significantly increased in range of motion for two of 19 tests, when compared to control. The description of the 3S training procedures revealed the omission of an important step - concentric contraction of the opposite muscle group together with light pressure from partner. Song and Garvie's method was:

> . . . method of increasing range of motion involves passively extending a muscle group close to the maximum and then, against a fixed resistance, isometrically contracting this muscle group while in the extended position for 5 to 7 second period. Following the contraction the muscle group is relaxed and once relaxed, further extension of the particular group is possible (p. 18).

Turner (1977) examined the effects of two flexibility methods, 3S and SS. The results showed flexibility training increased flexibility over weeks of training and flexibility decreased following post-training, when compared to control. There was no difference between the 3S and SS training methods during the course of the study. The 3S method employed accurately followed Holt's 3S standard procedures.

A comparison of the past research to the present study showed that only one study (Turner, 1977) followed Holt's

(1973) 3S training procedures. Since Turner did not examine within-session training, comparison was not made. Moreover, the present study found almost identical methods (e.g., 3S and CS training) did not produce the same significant effect. Based on this finding, comparison of similar methods in past research with dissimilar procedures was not warranted.

Theoretical Explanation

Herman Kabat, a medical doctor, originally drafted the proprioceptive neuromuscular facilitation (PNF) principles followed by techniques and methods of exercise to restore strength, endurance and range of motion in muscles of patients suffering from paralysis (e.g., poliomyelitis). The central goal of the PNF techniques was to maximally activate all functional motor units of the affected muscle during a voluntary muscular contraction (motion). The PNF techniques were defined by Kabat as being the following:

1) manual resistance,
2) stretch,
3) mass movement patterns,
4) reflexes, and
5) reversal of antagonist.

A number of PNF methods were combined in various ways for summation of facilitation effect. Kabat utilized the following PNF methods:

1) rhythmic stabilization,
2) isometric reversal of antagonist,

3) isometric reversal of antagonists with isometric contraction, and

4) quick reversal of antagonists.

Kabat pointed out that specific methods may prove more or less effective depending on the nature of the paralysis. For example, rhythmic stabilization is effective in patients with lower motor neuron lesions and lesions of the corticospinal tracts and basal ganglia. However, patients with cerebellar involvement were not affected by rhythmic stabilization.

In specific reference to the development of greatest flexibility, Kabat recommended that the therapist move the joint beyond the active range of motion and the subject perform a maximal isometric contraction (Kabat, 1952).

Following Kabat's work, Voss, D. and Knott, M., a team of physical therapists inspired by Kabat, extended the application and procedures of the PNF technique to patients suffering from non-paralytic dysfunction (e.g., arthritic patients) as well as paralytic dysfunction. Voss herself became a living example of the effectiveness of the PNF therapeutic exercise, as she successfully treated her own disability (Knott, 1973).

Voss (1967), with Knott's assistance standardized a method of therapeutic exercise derived from Kabat's earlier work. The new PNF method contained a number of factors:

1) manual contact,

2) vocal tones,

3) maximal contraction against resistance, and

4) passive and active extension.

Voss supported the use of the PNF technique(s) as they were based on well-grounded information from motor learning, neurophysiology, and motor behavior - research and clinical findings.

Subsequently, Holt appealed to individuals requiring increased range of motion to use the 3S flexibility training method. He suggests it is the only sport training method based on Herman Kabat's PNF (proprioceptive neuromuscular facilitation) therapeutic principles, and as a result is a firmly accepted neurophysiological fact.

Recently, researchers have begun to doubt Kabat's theoretical explanation of the advantages of PNF methods over other procedures which are entrenched in classical Sherringtonian views concerning successive inductive and reciprocal inhibition (Moore and Hutton, 1980). Furthermore, they stated, "the simplistic notion concerning reciprocal inhibition during muscle stretch should be discarded" (p. 327).

A review of selected literature suggested there are three major neural mechanisms (reciprocal inhibition, clasp knife reflex and successive induction) which may underlie the different flexibility training methods.

Reciprocal inhibition causes the antagonists (muscles opposing the contraction) to relax and lengthen, while the agonists (muscles contracting) shorten. For example, to move the forearm the biceps must contract, simultaneously the triceps must relax (Eccles, Fatt, and Landgren, 1956).

Clasp knife reflex causes the collapse of the muscles when various muscles or tendons are subjected to extensive force. For example, when a subject's arm becomes hyper-

extended during a football tackle, the muscles surrounding the shoulder joint collapse to help protect against injury (Rymer, Houk, and Crago, 1979).

Successive induction is the contraction of the muscles to be stretched (antagonists). Following successive induction, the muscles will be able to be further stretched, due to increased relaxation of the stretched muscle (Moore and Hutton, 1980) and a more forceful agonist contraction is facilitated (Holt, Kaplan, Okita, and Hoshika, 1969).

The present study revealed that the 3S training method was the most effective in increasing range of motion, across the ankle and shoulder joints, for within-session training. A comparison of almost identical methods (3S and CS training) on neural muscular mechanisms helped to explain the results.

The 3S and CS are identical, with one exception. In step four, 3S required both a concentric contraction with external force from a partner; while CS required only concentric contraction (Table 9). The 3S method activated both the reciprocal inhibition and clasp knife reflex, while CS method activated only reciprocal inhibition. The neural mechanisms underlying 3S training (reciprocal inhibition, clasp knife reflex and successive induction) suggested a possible explanation for the superiority of the 3S flexibility training method.

Within the scope of this study, it cannot be accurately determined which underlying neural mechanisms or other mechanisms caused the change in range of motion for each training method.

TABLE 9

NEURAL MECHANISMS UNDERLYING TRAINING METHODS

Method	Procedures	Neural Mechanisms	Effective Ankle	Effective Shoulder
3S	1. 2 sec concentric contraction	(RI)	*	*
	2. 6 sec isometric contraction	(SI)		
	3. 2 sec concentric contraction and external force	(RI & CK)		
2S	1. 2 sec passive extension	(CK)	*	
	2. 8 sec held passive extension	(CK)		
SS	1. 2 sec concentric contraction	(RI)		
	2. 8 sec isometric contraction	(RI)		
ES	1. 5 sec eccentric contraction	(SI)	*	
	2. 5 sec passive extension	(CK)		
BS	1. 1 sec concentric contraction (x 10)	(RI)		
CS	1. 2 sec concentric contraction	(RI)	*	
	2. 6 sec isometric contraction	(SI)		
	3. 2 sec concentric contraction	(RI)		

RI: reciprocal inhibition
CK: clasp knife reflex
SI: successive induction
*: significant ($p < 0.05$)

Future Research

It is recommended that future research be conducted on the mechanisms affecting range of motion. Once this is undertaken, a clearer understanding as to why particular methods enhance, retard, or do not affect flexibility will emerge. In addition, training methods superior to the 3S training technique used in the present investigation may also emerge.

Summary

The major finding and contributions of this study were as follows:

1) Scientific Stretching for Sport (3S) was the only training method that consistently increased range of motion on both ankle plantar flexion and shoulder extension for within-session training.
2) Six different standardized flexibility training methods (3S, 2S, SS, ES, BS, and CS) for ankle plantar flexion and shoulder extension were developed.
3) A modified Leighton Flexometer test method for ankle plantar flexion and shoulder extension was developed.

CHAPTER V

SUMMARY, CONCLUSIONS AND RECOMMENDATIONS

Summary

Even though flexibility has been recognized as being important in training and conditioning, and many years have been spent in examining the various joint flexibility responses to different stretching techniques, it is apparent from the literature that there is no universally accepted method of stretch training. This study investigated the effects of six different standardized stretch training methods on active flexibility for shoulder extension and ankle plantar flexion. The different training methods were: 1) 3S (concentric, isometric and concentric contractions with external force), 2) 2S (passive extension), 3) SS (concentric and isometric contractions), 4) ES (eccentric contraction and passive extension), 5) BS (concentric contraction), and 6) CS (concentric, isometric and concentric contractions). Forty-two female volunteer students ranging in age from 12 to 14 years were selected as subjects. Test-orientation and training-orientation sessions were followed by a six week flexibility training program and a two week post-training retention period. The modified Leighton Flexometer was utilized to test active range of motion for the ankle and shoulder joints.

The research design consisted of two 7 x 5 x 2 (treatment, week, and session) factorial designs with repeated measures

on the last two factors. Data was analyzed using two three-way analyses of variance (Greenhouse-Geisser test-on repeated measures) and post-hoc Scheffé test ($\alpha = 0.05$). The ankle and shoulder data were analyzed independently.

Conclusions

Within the scope of this study, the following conclusions were reached for ankle plantar flexion and shoulder extension.

Ankle plantar flexion. 1) The 3S, 2S, ES and CS flexibility training methods significantly increased ankle plantar flexion within-session training ($p < 0.05$).

2) The 3S training method increased ankle plantar flexion significantly more than CS training for within-session training ($p < 0.05$).

Shoulder extension. 1) The 3S training method significantly increased shoulder extension for within-session training ($p < 0.05$).

Comparison across joints. 1) The 3S training method consistently increased range of motion in both ankle plantar flexion and shoulder extension for within-session training ($p < 0.05$).

2) The SS and BS training methods consistently showed no increase in range of motion in both ankle plantar flexion and shoulder extension for within-session training ($p > 0.05$).

3) The changes in range of motion for ankle plantar flexion were not identical to changes in shoulder extension, over

the eight week period of time.

Recommendations

For further investigation, it is recommended that the present study be replicated using:

1) different age levels, ability levels, and sport and fitness groups (e.g., sedentary 25 to 30 year olds),
2) different numbers of repetitions, sets and frequencies,
3) different amounts of time for work and rest intervals,
4) an investigation on the immediate retention effects of flexibility (e.g., 30 seconds, one minute, and fifteen minutes), and
5) an examination of specific factors underlying all flexibility methods (e.g., external and internal forces).

BIBLIOGRAPHY

Adams, J.C. Outline of orthopedics. London: Churchill Livingstone, 1971.

Atha, J. and Wheatley, D.W. The mobilizing effects of repeated measurements on hip flexion. British Journal of Sport Medicine, 1976, 10, 22-24.

Barnsley, R.H. and Rabinovitch, M.S. Handedness: Proficiency versus stated preference. Perceptual and Motor Skills, 1970, 30, 342-362.

Broer, M.R. and Galles, N.R.G. Importance of relationship between various body measurements in performance of the toe-touch test. Research Quarterly, 1959, 29, 253-263.

Bushey, S.R. Relationship of modern dance performance to agility, balance, flexibility, power, and strength. Research Quarterly, 1966, 37, 313-316.

Carlile, F. The effects of preliminary passive warming on swimming. Research Quarterly, 1956, 27, 143-151.

Chapman, E.A., de Vries, H.A. and Swezey, R. Joint stiffness: effects of exercise on young and old men. Journal of Gerontology, 1972, 27, 218-221.

Cherry, D.B. Review of physical therapy alternatives for reducing muscle contracture. Physical Therapy, 1980, 60, 877-881.

Clarke, H. Joint and body range of movement. Physical Fitness Research Digest, 1975, 5, 1-22.

Cook, T.D. and Campbell, D.T. Quasi-experimentation: Design and analysis issues for field settings. Chicago: Rand McNally, 1979.

Cooper, D.L. and Fair, J. Stretching exercise for flexibility. The Physician and Sports Medicine, 1977, 5, 114-116.

Corbin, C.B. and Noble, L. Flexibility a major component of physical fitness. Journal of Health, Physical Education and Recreation, 1980, 51, 23-24 and 57-60.

Cotten, D.J. A comparison of selected trunk flexibility tests. American Corrective Therapy Journal, 1972, 26, 24-26.

Cureton, T.K. Flexibility as an aspect of physical fitness. Research Quarterly, 1941, 12, 381-390.

Davies, E.A. Relationship between selected postural divergencies and motor ability. Research Quarterly, 1957, 28, 1-4.

Denk, G.M. The changes occurring in strength and flexibility during a competitive gymnastics season involving high school boys. Unpublished Master's Thesis, University of Kansas, 1971.

de Vries, H.A. Prevention of muscular distress after exercise. Research Quarterly, 1961, 32, 177-185.

de Vries, H.A. Evaluation of static stretching procedures for improvement of flexibility. Research Quarterly, 1962, 33, 222-229 and 656.

de Vries, H.A. The "looseness" factor in speed and oxygen consumption of an anaerobic 100 yard dash. Research Quarterly, 1963, 34, 305-313.

de Vries, H.A. Physiology of exercise. Iowa: WM. C. Brown Co., 1976.

Dickinson, R.V. The specificity of flexibility. Research Quarterly, 1968, 39, 792-794.

Dinkheller, A.L.B. Factors affecting flexibility. Unpublished Master's Thesis, University of Iowa, 1969.

Downie, P.D. A study of the relationship between flexibility measures and chronological ages of six to ten year old girls. Unpublished Master's Thesis, University of Oregon, 1965.

Downie, P.D. A study of the flexibility characteristics of ten, eleven, twelve, thirteen and fourteen year old girls. Unpublished Doctoral Dissertation, University of Oregon, 1970.

Eccles, J.C., Fatt, P. and Landgren, S. The central pathway for the direct inhibitory action of impulses in the largest afferent nerve fibres to muscle. Journal of Neurophysiology, 1956, 126, 524-562.

Fieldman, H. Effects of selected extensibility exercises on the flexibility of the hip joint. Research Quarterly, 37, 1966, 326-331.

Francis, C.C. Introduction to human anatomy. Saint Louis: The C.V. Mosby Company, 1968.

Harris, M.L. A factor analytic study of flexibility. Research Quarterly, 1969, 40, 62-70.

Hartley, S. A comparison of six methods of stretch on active range of hip flexion. Unpublished Master's Thesis, University of British Columbia, 1976.

Harvey, V.P. and Scott, G.D. Reliability of a measure of forward flexibility and its relation to physical dimensions of college women. Research Quarterly, 1967, 38, 28-33.

Hogg, J.M. Flexibility and its importance for the competitive swimmer. Unpublished paper presented at the 9th World Swimming Clinic, San Diego, California, 1977.

Hogg, J.M. Flexibility training: Its importance for the competitive swimmer. Katimavik, 1978, 5, 1-6.

Holt, L.E., Kaplan, H.M., Okita, T. and Hoshiko, M. The influence of antagonistics contraction and head position on the responses of agonistic muscles. Archives of Physical Medicine and Rehabilitation, 1969, 50, 283-291.

Holt, L.E., Travis, T.M. and Okita, T. Comparative study of three stretching techniques. Perceptual and Motor Skills, 1970, 31, 611-616.

Holt, L.E. Scientific Stretching for Sport. Halifax: Dalhousie University, 1973.

Hupprich, F.L. and Sigerseth, P.O. The specificity of flexibility in girls. Research Quarterly, 1949, 21, 25-33.

Jacobs, M. Neurophysiological implications of slow active stretching. American Corrective Therapy Journal, 1976, 30, 151-154.

Jesse, J.P. Misuse of strength development programs in athletic training. The Physician and Sports Medicine, 1979, 7, 46-52.

Johns, R.J. and Wright, V. Relative importance of various tissues in joint stiffness. Journal of Applied Physiology, 1962, 17, 824-828.

Kabat, H. Studies on neuromuscular dysfunction XV: The role of central facilitation in restoration of motor function in paralysis. Archives of Physical Medicine and Rehabilitation, 1952, 33, 521-533.

Keppel, G. Design and analysis. A researcher's handbook. New Jersey: Prentice-Hall, Inc., 1973.

Kingsley, D.B. Flexibility changes resulting from participation in tumbling. Unpublished Master's Thesis, University of Oregon, 1952.

Kirchner, G. and Glines, D. Comparative analysis of Eugene, Oregon, elementary school children using the Kraus-Weber test of minimum muscular fitness. Research Quarterly, 1957, 28, 16-25.

Knott, M. In the groove. Physical Therapy, 1973, 53, 365-372.

Kraus, H. and Hirschland, R.P. Minimum muscular fitness tests in school children. Research Quarterly, 1954, 25, 178-188.

Last, R.J. Anatomy regional and applied. New York: Churchill Livingstone, 1978.

Laubach, L.L. and McConville, J.T. Relationships between flexibility, anthropometry and the somatotype of college men. Research Quarterly, 1966(a), 37, 241-251.

Laubach, L.L. and J.T. McConville. Muscle strength, flexibility, and body size of adult males. Research Quarterly, 1966(b), 37, 384-392.

Leighton, J.R. An investigation of the flexibility characteristics of three age groups and five specialized skill groups among males twelve, fourteen, sixteen years of age and college age athletes. Unpublished Doctoral Dissertation, University of Oregon, 1954.

Leighton, J.R. Flexibility characteristics of males ten to eighteen years of age. Archives of Physical Medicine and Rehabilitation, 1956, 37, 494-499.

Leighton, J.R. Flexibility characteristics of four specialized skill groups of college athletes. Archives of Physical Medicine and Rehabilitation, 1957, 32, 24-28.

Leighton, J.R. On the significance of flexibility for physical educators. Journal of Health, Physical Education and Recreation, 1960, 31, 27-28.

Leighton, J.R. The Leighton flexometer and flexibility test. Journal of the Association for Physical and Mental Rehabilitation, 1966, 20, 86-93.

Lemiere, O. Flexibility of shot putters and discus throwers. Unpublished Master's Thesis, University of Oregon, 1952.

Logan, G.A. and Egstrom, G.H. Effects of slow and fast stretching on the sacro-femoral angle. Journal of the Association for Physical and Mental Rehabilitation, 1961, 15, 85-89.

Massey, B.H. and Chaudet, N.L. Effects of systematic, heavy resistance exercise on range of joint movement in young male adults. Research Quarterly, 1956, 27, 41-51.

Mathews, D.K., Shaw, V. and Bohnen, M. Hip flexibility of college women as related to length of body segments. Research Quarterly, 1957, 28, 352-356.

McCue, B.F. Flexibility measurements of college women. Research Quarterly, 1952, 24, 316-324.

McGavin, J.C. A comparison of constant and variable resistance training upon selected aspects of shoulder dynamics. Unpublished Master's Thesis, The University of Alberta, 1980.

Medeiros, J.M., Smidt, G.L., Burmeister, L.F. and Soderberg, G.L. The influence of isometric exercise and passive stretch on hip joint motion. Physical Therapy, 1974, 40, 518-523.

Moore, M.A. and Hutton, R.S. Electro-myographic investigation of muscle strength techniques. Medicine and Science in Sports and Exercise, 1980, 12, 323-329.

Phillips, M., Bookwalter, C., Denman, C., McAuley, J., Sherwin, H., Summers, D., and Yeakel, H. Analysis of results from the Kraus-Weber test of minimum muscular fitness in children. Research Quarterly, 1955, 26, 314-323.

Rochelle, R.H., Skubic, V., and Michael, E.D. Performance as affected by incentive and preliminary warmup. Research Quarterly, 1960, 31, 499-504.

Ruhl, P.F. Changes in strength, flexibility, balance, movement time and dance techniques occurring during University of Kansas gymnastics season. Unpublished Master's Thesis, University of Kansas, 1972.

Rymer, W.Z., Houk, J.C. and Crago, P.E. Mechanisms of the clasp-knife reflex studied in an animal model. Exp. Brain Res., 1979, 37, 93-113.

Schultz, P. Flexibility: Day of the static stretch. The Physician and Sports Medicine, 1979, 7, 109-117.

Scott, M.G. and French, E. Measurement and evaluation in physical education. Dubuque: Iowa: Wm. C. Brown, 1959.

Sigerseth, P.O. and Haliski, C.C. The flexibility of football players. Research Quarterly, 1950, 21, 394-398.

Song, T.M.K. and Garvie, G.T. Wrestling with flexibility. C.A.H.P.E.R. Journal, 1976, 43, 18-26.

Song, T.M.K. "Flexibility of ice hockey players and comparison with other groups." In Terauds, J. and Gros, H.J. (eds.) Science in skiing, skating and hockey. California: Academic Publishers, 1979, 117-125.

Song, T.M.K. and Reid, R. "Relationship of lower limb flexibility, strength and anthropometric measures to skating speed." In Terauds, J. and Gros, H.J. (eds.) Science in skiing, skating and hockey. California: Academic Publishers, 1979, 83-98.

Stein, R.B. Nerve and muscle membranes, cells and systems. London: Plenum Press, 1980.

Tanigawa, M.C. Comparison of the hold-relax procedure and passive mobilization on increasing muscle length. Physical Therapy, 1972, 52, 725-735.

Trombly, C.A. and Scott, A.D. Occupational therapy for physical dysfunction. London: Williams and Wilkins, 1976.

Tucker, N.M. Changes in the range of movement of the arm following isotonic and isometric work to strengthen the musculature around the scapulohumeral joint. Unpublished Master's Thesis, University of Oregon, 1963.

Turner, A.A. The effects of two training methods of flexibility. Unpublished Master's Thesis, Lakehead University, 1977.

Turner, A., and Singh, M. Study of traditional and modified joint flexibility tests. Proceedings of the SIPT Conference held in Red Deer, Alberta, 1981.

Tyrance, H.J. Relationships of extreme body types to ranges of flexibility. Research Quarterly, 1958, 29, 349-359.

Voss, D.E. Proprioceptive neuromuscular facilitation. American Journal of Physiology and Medicine, 1967, 46, 838-898.

Wear, C.L. Relationship of flexibility measurements to length of body segments. Research Quarterly, 1963, 34, 234-238.

Weber, S. and Kraus, H. Passive and active stretching of muscles: Spring stretch and control group. The Physical Therapy Review, 1949, 29, 407-410.

Wells, K.F. and Dillon, E.K. The sit and reach - a test of back and leg flexibility. Research Quarterly, 1952, 23, 115-118.

Wickstrom, R.L. The effect of low-resistance, high repetition progressive resistance exercise upon selected measures of strength and flexibility. Journal of the Association for Physical and Mental Rehabilitation, 1960, 14, 161-162 and 173.

Wiechec, F.J. and Krusen, F.H. A new method of joint measurement and a review of the literature. American Journal of Surgery, 1939, 43, 659-668.

Winer, B.J. Statistical principles in experimental design. New York: McGraw-Hill Book Company, 1971.

APPENDIX A

FLEXIBILITY TRAINING METHODS

SCIENTIFIC STRETCH FOR SPORT TRAINING

(3S)

Definition

This is a method of increasing flexibility whereby an isometric contraction of the muscles to be stretched (muscles in a lengthened position to start) is followed by a concentric contraction of the opposite muscle group, together with light force from a partner (external force), performed in a progressively more strenuous series until the range of movement cannot be increased without severe pain (Holt, 1973, pp. 5-7).

Explanation

A maximal range of motion is attained by actively and passively flexing or extending a segment nearest to the joint being trained, in an arc, as far as possible. Repetitions of the exercise are performed, without severe pain. One repetition consists of:

a) two second concentric contraction,
b) six second isometric contraction, and
c) two second concentric contraction, with light force from a partner.

Subject repeats the above exercise (rest optional). The contractions performed by the exercise subject must be gradually, not explosively, increased in effort.

(3S)

Shoulder Extension Exercise (Stretching Shoulder Flexors)

Step 1. Subject (S) assumes a prone position on the floor, legs and back straight, arms down at sides, with hands supinated. Partner (P) stands in front of S, in an anatomical position.



(3S)

Step 2. Subject's (S's) training arm is extended, in an arc, at the shoulder joint as far as possible, with hand in a neutral position, for a vocal count of two (seconds).



(3S)

Step 3. Partner (P) flexes at the hips, holding S's arm proximal to the wrist. S attempts to flex the arm, while keeping the elbows straight. P resists S's movement. P holds S's position for a vocal count of six (seconds) to produce an isometric contraction.



(3S)

Step 4. Subject (S) again fully extends the arm for a vocal count of two (seconds) keeping the elbow straight (1st), together with light force from P (2nd), so that S attains the maximum extended range of motion in the shoulder joint, without severe pain.



(3S)

Step 5. Subject (S) relaxes for a vocal count of fifteen (seconds) (step 1), then repeats steps two, three, and four. The exercise is performed three consecutive times.

(3S)

Ankle Plantar Flexion Exercise (Stretching Ankle Flexors)

Step 1. Subject (S) assumes a sitting position on the floor, legs and back straight, arms at sides, with hands pronated. Partner (P) kneels in front of S, in a partial anatomical position.



(3S)

Step 2. Subject's (S's) training feet are plantar flexed, in an arc, at the ankle joint, as far as possible, for a vocal count of two (seconds).



(3S)

Step 3. Partner (P) flexes at the hips, holding S's feet distal to the ankles. S attempts to dorsi-flex the feet, while keeping the knees straight. P resists S's movement. P holds S's position for a vocal count of six (seconds) to produce an isometric contraction.



(3S)

Step 4. Subject (S) again fully plantar flexes the feet for a vocal count of two (seconds) keeping the knees straight (1st), together with light force from P (2nd), so that S attains the maximum flexed range of motion in the ankle joint, without severe pain.



(3S)

Step 5. Subject (S) relaxes for a vocal count of fifteen (seconds) (step 1), then repeats steps two, three, and four. The exercise is performed three consecutive times.

STATIC STRETCH TRAINING

(2S)

Definition

This is a method of increasing flexibility whereby an external force (i.e., hands pulling the segment) is applied in a progressively more forceful manner, until the range of movement cannot be increased, then held for a period of time, without severe pain (de Vries, 1976, pp. 436-437).

Explanation

A maximal range of motion is attained by passively flexing or extending a segment nearest to the joint being trained, in an arc, as far as possible. Repetitions of the exercise are performed without severe pain. One repetition consists of:

a) two second passive extension, and

b) eight second held passive extension.

Subject relaxes for fifteen seconds before repeating each exercise. The external force applied to extend the segment is gradually, not explosively, increased in effort.

(2S)

Shoulder Extension Exercise (Stretching Shoulder Flexors)

Step 1. Subject (S) assumes a prone position on the floor, legs and back straight, arms down at sides, with hands supinated. Partner (P) stands in front of S, in an anatomical position.



(2S)

Step 2. Partner (P) flexes at the hips, holding S's arm proximal to the wrist. S relaxes the shoulder, keeping the elbow extended, hand in a neutral position, while P passively extends the arm for a vocal count of two (seconds).



(2S)

Step 3. Partner (P) fully extends the arm, without severe pain, and this position is held for a vocal count of eight (seconds).



(2S)

Step 4. Subject (S) relaxes for a vocal count of fifteen (seconds) (step 1), then repeats step two and three. The exercise is performed three consecutive times.

(2S)

Ankle Plantar Flexion Exercise (Stretching Dorsal Flexors)

Step 1. Subject (S) assumes a sitting position on the floor, legs and back straight, arms at sides, with hands pronated. Partner (P) kneels in front of S, in a partial anatomical position.



(2S)

Step 2. Partner (P) flexes at the hips, holding S's feet distal to the ankle. S relaxes the ankle, keeping the knees extended, while P passively extends the feet for a vocal count of two (seconds).



(2S)

Step 3. Partner (P) fully extends the feet, without severe pain, and this position is held for a vocal count of eight (seconds).



(2S)

Step 4. Subject (S) relaxes for a vocal count of fifteen (seconds) (step 1), then repeats steps two and three. The exercise is performed three consecutive times.

SLOW ACTIVE STRETCH TRAINING

(SS)

Definition

This is a method of increasing flexibility by slow active concentric contraction of the agonist muscles, without additional assistance from an external force (i.e., hands pulling the segment to a greater range of motion), while relaxing the antagonist muscle group (the muscles to be stretched) (Jacobs, 1976, pp. 151-152).

Explanation

A maximal range of motion is attained by actively flexing or extending a segment nearest to the joint being trained, in an arc, as far as possible. Repetitions of the exercise are performed, without severe pain. One repetition consists of:

a) two second concentric contraction, and

b) eight second isometric contraction.

Subject relaxes for fifteen seconds before repeating each exercise. The contractions performed by the subject must be gradually, not explosively, increased in effort.

(SS)

Shoulder Extension Exercise (Stretching Shoulder Flexors)

Step 1. Subject (S) assumes a prone position on the floor, legs and back straight, arms down at sides, with hands supinated.



(SS)

Step 2. Subject's (S's) training arm is extended, in an arc, at the shoulder joint, as far as possible for a vocal count of two (seconds), then this position is held for a vocal count of eight (seconds). The subject's elbow is extended, with hand in a neutral position.



(SS)

Step 3. Subject (S) relaxes for a vocal count of fifteen (seconds) (step 1), then repeats step two. The exercise is performed three consecutive times.

(SS)

Ankle Plantar Flexion Exercise (Stretching Dorsal Flexors)

Step 1. Subject (S) assumes a sitting position on the floor, legs and back straight, arms at sides, with hands pronated.



(SS)

Step 2. Subject's (S's) training feet are plantar flexed in an arc, at the ankle joint, as far as possible, for a vocal count of two (seconds), then this position is held for a vocal count of eight (seconds).



(SS)

Step 3. Subject (S) relaxes for a vocal count of fifteen (seconds) (step 1), then repeats step two. The exercise is performed three consecutive times.

ECCENTRIC STRETCH TRAINING

(ES)

Definition

This is a method of increasing flexibility whereby a slow active eccentric contraction of the muscles to be stretched (muscles in a contracted position to start) is followed by light force from a partner (external force), while the subject relaxes both agonist and antagonist muscles around the joint.

Explanation

A maximal range of motion is attained by actively flexing or extending a segment nearest to the joint being trained, in an arc, as far as possible. Repetitions of the exercise are performed, without severe pain. One repetition consists of:

a) five second eccentric contraction, and

b) five second passive extension.

Subject relaxes for fifteen seconds before repeating each exercise. The contraction and passive extension performed by the subject must be gradually, not explosively, increased in effort.

(ES)

Shoulder Extension Exercise (Stretching Shoulder Flexors)

Step 1. Subject (S) assumes a prone position on the floor, legs and back straight, arms at sides, with hands supinated. Partner (P) stands in front of S, in an anatomical position.



(ES)

Step 2. Partner (P) flexes at the hips, holding S's arm proximal to the wrist.



(ES)

Step 3. Subject (S) attempts to flex the arm, while keeping the elbow straight, with hand in a neutral position. P, with greater force, extends S's arm, in an arc, as far as possible, without severe pain, for a vocal count of five (seconds).



(ES)

Step 4. Subject (S) relaxes, while P passively extends S's arm, as far as possible, keeping the elbow straight, with hand in a neutral position, for a vocal count of five (seconds).



(ES)

Step 5. Subject (S) relaxes for a vocal count of fifteen (seconds), then repeats steps two, three and four. The exercise is performed three consecutive times.

(ES)

Ankle Extension Exercise (Stretching Ankle Flexors)

Step 1. Subject (S) assumes a sitting position on the floor, legs and back straight, arms at sides, with hands pronated. Partner (P) kneels in front of S, in a partial anatomical position.



(ES)

Step 2. Partner (P) flexes at the hips, holding S's training feet distal to the ankles.



(ES)

Step 3. Subject (S) attempts to dorsi-flex the feet, while keeping the knees straight. P, with greater force, extends S's feet, in an arc, as far as possible, without severe pain for a vocal count of five (seconds).



(ES)

Step 4. Subject (S) relaxes, while P passively extends S's feet, as far as possible, keeping the knees straight, for a vocal count of five (seconds).



(ES)

Step 5. Subject (S) relaxes for a vocal count of fifteen (seconds) (step 1), then repeats steps two, three and four. The exercise is performed three consecutive times.

BALLISTIC STRETCH TRAINING

(BS)

Definition

This is a method of increasing flexibility whereby particular emphasis is placed on training each individual to make all movements in a relaxed rhythmic manner, but to put special effort toward carrying the movement to the limit of the subject's range of motion (McCue, 1953, p. 322).

Explanation

A maximal range of motion is attained by actively flexing or extending, in a relaxed rhythmic manner, a segment nearest to the joint being trained, in an arc, as far as possible. Repetitions of the exercise are performed, without severe pain. One repetition consists of one second concentric contractions. Subjects will perform 10 consecutive repetitions, then relax for fifteen seconds before repeating each exercise. The series of ten concentric contractions is performed in a rhythmic manner, explosive in effort.

(BS)

Shoulder Extension Exercise (Stretching Shculder Flexors)

Step 1. Subject (S) assumes a prone position on the floor, legs and back straight, arms at sides, with hands supinated.



(BS)

Step 2. In a relaxed rhythmic manner, the training arm is extended in an arc, at the shoulder joint, as far as possible, without holding the position, for ten consecutive repetitions. The subject's elbow is extended, with hand in a neutral position.



(BS)

Step 3. Subject (S) relaxes for a vocal count of fifteen (seconds) (step 1), then repeats step two. The exercise is performed three consecutive times.

(BS)

Shoulder Extension Exercise (Stretching Shoulder Flexors)

Step 1. Subject (S) assumes a sitting position on the floor, legs and back straight, arms at sides, with hands pronated.



(BS)

Step 2. In a relaxed rhythmic manner the training feet are plantar flexed in an arc, at the ankle joint, as far as possible, without holding the position, for ten consecutive repetitions.



(BS)

Step 3. Subject (S) relaxes for a vocal count of fifteen (seconds) (step 1), then repeats step two. The exercise is performed three consecutive times.

CONTRACT RELAX STRETCH TRAINING

(CS)

Definition

This is a method of increasing flexibility whereby an isometric contraction of the muscles to be stretched (muscles in a lengthened position to start) is followed by a concentric contraction of the opposite muscle group (Moore and Hutton, 1980, p. 323).

Explanation

A maximal range of motion is attained by actively flexing or extending a segment nearest to the joint being trained, in an arc, as far as possible. Repetitions of the exercise are performed, without severe pain. One repetition consists of:

a) two second concentric contraction,

b) six second isometric contraction, and

c) two second concentric contraction.

Subject immediately repeats the above exercise (rest optional). The isometric contraction performed by the exercise subject must be gradually, not explosively, increased in effort.

(CS)

Shoulder Extension (Stretching Shoulder Flexors)

Step 1. Subject (S) assumes a prone position on the floor, legs and back straight, arms at sides, with hands supinated. Partner (P) stands in front of S, in an anatomical position.



(CS)

Step 2. Subject's (S's) training arm is extended, in an arc, at the shoulder joint as far as possible, with hand in a neutral position, for a vocal count of two (seconds).



(CS)

Step 3. Partner (P) flexes at the hips, holding S's arm proximal to the wrist. S attempts to flex the arm, while keeping the elbows extended. P resists S's movement. P holds S's position for a vocal count of six (seconds) to produce an isometric contraction.



(CS)

Step 4. Subject (S) again fully extends the arm for a vocal count of two (seconds) keeping the elbow straight so that S attains the maximum extended range of motion in the shoulder joint, without severe pain.



(CS)

Step 5. Subject (S) relaxes for a vocal count of fifteen (seconds) (step 1), then repeats steps two, three and four.

(CS)

Ankle Plantar Flexion Exercise (Stretching Dorsal Flexors)

Step 1. Subject (S) assumes a sitting position on the floor, legs and back straight, arms at sides, with hands pronated. Partner (P) kneels in front of S, in partial anatomical position.



(CS)

Step 2. Subject's (S's) training feet are plantar flexed, in an arc, at the ankle joint, as far as possible, for a vocal count of two (seconds).



(CS)

Step 3. Partner (P) flexes at the hips, holding S's feet distal to the ankles. S attempts to dorsi-flex the feet, while keeping the knees extended. P resists S's movement. P holds S's position for a vocal count of six (seconds) to produce an isometric contraction.



(CS)

Step 4. Subject (S) again fully plantar flexes the feet for a vocal count of two (seconds) keeping the knees straight so that S attains the maximum flexed range of motion in the ankle joint, without severe pain.



(CS)

Step 5. Subject (S) relaxes for a vocal count of fifteen (seconds) (step 1), then repeats steps two, three and four. The exercise is performed three consecutive times.

APPENDIX B

FLEXIBILITY TESTS AND APPARATUS

FLEXIBILITY TESTS

Shoulder Extension Test (Stretching Shoulder Flexors)

Step 1. Subject (S) assumes a prone position on the bench, legs and back straight, arms at sides, with hands in the neutral position.



Step 2. The flexometer is strapped on the right dominant, lateral side of the arm, proximal to the elbow joint, with test shoulder beyond the bench. The portable wall is placed medial to the test arm.



Step 3. Subject's (S's) test arm is extended, in an arc, at the shoulder joint, with hand in a neutral position, for a vocal count of two (seconds). The palm of the hand slides lightly against the wall, as far as possible, and this position is held for a vocal count of three (seconds). The highest measurement is read and recorded.



Step 4. Subject (S) relaxes for a vocal count of fifteen (seconds) (step 1), then repeats steps two and three. The test is performed three consecutive times.



Ankle Plantar Flexion Test (Stretching Dorsal Flexors)

Step 1. Subject (S) assumes a sitting position on the bench, back straight, arms at sides.. The test leg is flexed at the knee by placing a cylinder posterior to the knee joint, with the inactive foot at the side of the bench.



Step 2. Subject (S) is strapped to the bench, proximal to the knee joint. The flexometer is attached to the left dominant medial arch, with test ankle beyond the bench.



Step 3. Subject's (S's) test foot is plantar flexed, in an arc, at the ankle joint, for a vocal count of two (seconds). The ankle is plantar flexed, as far as possible, and this position is held for a vocal count of three (seconds). The highest measurement is read and recorded.



Step 4. Subject (S) relaxes for a vocal count of fifteen (seconds) (step 1), then repeats steps two and three. The test is performed three consecutive times.





Figure A. Test cylinder for the ankle plantar flexion test.



Figure B. Test strap for the shoulder extension test.



Figure C. The Leighton Flexometer.

APPENDIX C

CORRESPONDENCE

Lakewood Intermediate School
THIRD STREET N. · KENORA, ONTARIO P9N 3Y5 · PHONE 468-3131

November 6, 1981.

To Whom It May Conern,

Re: Flexibility Training Programme

Please be advised that Mr. Allan Turner has been conducting research on a Flexibility Training Programme for the months of September through December at Lakewood Intermediate School. Specifically, Allan has worked with two classes of girls physical education to compile the results which will form part of his Doctorate research.

We were pleased to have Allan in the school as he provided our students the opportunity to witness, first-hand, research being conducted; and, on several occasions, assisted us as a supply teacher. We are indeed grateful that Allan chose Lakewood for his research project and wish him every success in his future endeavours.

Sincerely,

R.C. Harvey,
Principal.

RCH/pb

112 - 8604 - 103 Street
Edmonton, Alberta
T6E 4B6

February 14, 1981

Mr. R. Harvey
Principal
Lakewood Intermediate School

Dear Mr. Harvey,

The purpose of this letter is to request permission to conduct a research study in Lakewood Intermediate School, from September to October, 1981.

The aim of this study will be to examine the effects of six different stretch training methods on active joint flexibility. Subjects will be 42 female junior high school students and the design will consist of six different groups with six students in each group. The students will be required to perform one exercise on the ankle and one on the shoulder joint at the beginning of each physical education class. The subjects will be tested before training, after two, four and six weeks of training, and two weeks following no flexibility training (retention test).

Since this program will be partial requirement for my doctoral thesis, I would sincerely appreciate your assistance regarding this request. Should further information be required, I will be pleased to provide it (1-403-433-0956).

Sincerely,

A. Allan Turner

Andrew Allan Turner
Prov. Ph.D. Candidate

INFORMATION SHEET

1) NAME: _____

2) ADDRESS: _____

_____

3) PHONE: _____

4) AGE: _____

5) GRADE: _____

6) HAND DOMINANCE: R L (circle one)

7) PREVIOUS INJURY/DISEASE TO ANKLE OR SHOULDER: YES NO

8) IF YES, EXPLAIN WHAT AND WHERE: _____

_____

_____

_____

9) SURGERY ON JOINTS: YES NO

10) LIST ACTIVITIES YOU WILL BE PARTICIPATING IN DURING THE NEXT EIGHT WEEKS.

1) _____ 6) _____

2) _____ 7) _____

3) _____ 8) _____

4) _____ 9) _____

5) _____ 10) _____

APPENDIX D

RAW DATA

TABLE A

RAW SCORES (IN DEGREES) OF ANKLE JOINT FLEXIBILITY FOR EACH SUBJECT

#	SAB	C_1	C_2
1	1 1 1	112.00	117.00
2	1 1 2	108.00	114.00
3	1 1 3	111.00	116.00
4	1 1 4	116.00	121.00
5	1 1 5	113.00	117.00
6	2 1 1	090.00	094.00
7	2 1 2	091.00	094.00
8	2 1 3	093.00	094.00
9	2 1 4	100.00	100.00
10	2 1 5	093.00	093.00
11	3 1 1	108.00	108.00
12	3 1 2	102.00	108.00
13	3 1 3	103.00	108.00
14	3 1 4	104.00	107.00
15	3 1 5	098.00	104.00
16	4 1 1	108.00	112.00
17	4 1 2	108.00	108.00
18	4 1 3	108.00	110.00
19	4 1 4	116.00	116.00
20	4 1 5	110.00	114.00
21	5 1 1	102.00	108.00
22	5 1 2	108.00	108.00
23	5 1 3	106.00	106.00
24	5 1 4	107.00	111.00
25	5 1 5	104.00	108.00
26	6 1 1	128.00	133.00
27	6 1 2	124.00	127.00
28	6 1 3	133.00	136.00
29	6 1 4	138.00	138.00
30	6 1 5	133.00	136.00
31	1 2 1	107.00	112.00
32	1 2 2	102.00	108.00
33	1 2 3	106.00	106.00
34	1 2 4	111.00	111.00
35	1 2 5	101.00	112.00
36	2 2 1	108.00	111.00
37	2 2 2	108.00	113.00
38	2 2 3	107.00	110.00
39	2 2 4	114.00	117.00
40	2 2 5	111.00	114.00
41	3 2 1	098.00	103.00
42	3 2 2	100.00	101.00
43	3 2 3	102.00	103.00
44	3 2 4	106.00	107.00
45	3 2 5	100.00	103.00
46	4 2 1	102.00	106.00
47	4 2 2	103.00	108.00
48	4 2 3	103.00	107.00
49	4 2 4	111.00	113.00
50	4 2 5	104.00	107.00
51	5 2 1	108.00	111.00
52	5 2 2	110.00	114.00
53	5 2 3	108.00	112.00
54	5 2 4	113.00	116.00
55	5 2 5	110.00	113.00
56	6 2 1	101.00	101.00
57	6 2 2	102.00	104.00
58	6 2 3	102.00	105.00
59	6 2 4	105.00	113.00
60	6 2 5	100.00	103.00
61	1 3 1	101.00	101.00
62	1 3 2	107.00	108.00
63	1 3 3	108.00	108.00
64	1 3 4	110.00	110.00
65	1 3 5	110.00	108.00
66	2 3 1	104.00	106.00
67	2 3 2	101.00	103.00
68	2 3 3	102.00	103.00
69	2 3 4	111.00	111.00
70	2 3 5	108.00	108.00
71	3 3 1	116.00	117.00
72	3 3 2	117.00	118.00
73	3 3 3	120.00	119.00
74	3 3 4	125.00	124.00
75	3 3 5	121.00	120.00
76	4 3 1	110.00	107.00
77	4 3 2	108.00	107.00
78	4 3 3	106.00	104.00
79	4 3 4	111.00	112.00
80	4 3 5	110.00	108.00
81	5 3 1	088.00	093.00
82	5 3 2	088.00	088.00
83	5 3 3	086.00	084.00
84	5 3 4	088.00	084.00
85	5 3 5	093.00	090.00
86	6 3 1	097.00	097.00
87	6 3 2	098.00	098.00
88	6 3 3	100.00	097.00
89	6 3 4	098.00	101.00
90	6 3 5	097.00	098.00
91	1 4 1	111.00	113.00
92	1 4 2	108.00	111.00
93	1 4 3	108.00	113.00
94	1 4 4	118.00	120.00
95	1 4 5	111.00	118.00
96	2 4 1	108.00	111.00
97	2 4 2	108.00	108.00
98	2 4 3	110.00	113.00
99	2 4 4	112.00	118.00
100	2 4 5	107.00	111.00
101	3 4 1	088.00	093.00
102	3 4 2	090.00	088.00
103	3 4 3	090.00	094.00
104	3 4 4	094.00	098.00
105	3 4 5	093.00	097.00
106	4 4 1	098.00	098.00
107	4 4 2	098.00	098.00
108	4 4 3	097.00	098.00
109	4 4 4	098.00	100.00
110	4 4 5	098.00	100.00
111	5 4 1	108.00	108.00
112	5 4 2	108.00	108.00
113	5 4 3	108.00	108.00
114	5 4 4	112.00	114.00
115	5 4 5	108.00	108.00
116	6 4 1	116.00	116.00
117	6 4 2	116.00	117.00
118	6 4 3	116.00	117.00
119	6 4 4	116.00	116.00
120	6 4 5	116.00	116.00

Symbols

- # Number
- S Subject
- A Treatment
- B Week
- C Session

Treatment Conditions

- A_1 3S training (n = 6)
- A_2 2S training (n = 6)
- A_3 SS training (n = 6)
- A_4 ES training (n = 6)
- A_5 BS training (n = 6)
- A_6 CS training (n = 6)
- A_7 Control (n = 6)

Week in Experiment

- B_1 pre-training
- B_2 two-week training
- B_3 four-week training
- B_4 six-week training
- B_5 two-week retention

Session

- C_1 pre-session training
- C_2 post-session training

TABLE B

RAW SCORES (IN DEGREES) OF ANKLE JOINT FLEXIBILITY FOR EACH SUBJECT

#	SAB	C_1	C_2
121	1 5 1	107.00	105.50
122	1 5 2	111.50	110.50
123	1 5 3	107.50	107.00
124	1 5 4	114.50	114.50
125	1 5 5	108.50	110.50
126	2 5 1	094.50	095.00
127	2 5 2	095.00	094.00
128	2 5 3	093.50	093.50
129	2 5 4	101.50	099.50
130	2 5 5	094.50	095.50
131	3 5 1	099.00	103.50
132	3 5 2	106.50	103.50
133	3 5 3	107.00	107.00
134	3 5 4	113.50	113.50
135	3 5 5	108.50	108.50
136	4 5 1	112.50	112.00
137	4 5 2	112.00	111.50
138	4 5 3	112.00	116.00
139	4 5 4	112.50	111.50
140	4 5 5	108.50	108.50
141	5 5 1	101.50	102.50
142	5 5 2	102.00	103.00
143	5 5 3	103.50	103.00
144	5 5 4	108.00	107.50
145	5 5 5	100.50	101.50
146	6 5 1	113.50	116.00
147	6 5 2	115.00	116.50
148	6 5 3	118.00	118.00
149	6 5 4	117.00	121.00
150	6 5 5	110.50	110.00
151	1 6 1	110.50	110.50
152	1 6 2	110.00	111.50
153	1 6 3	107.50	108.50
154	1 6 4	114.50	113.50
155	1 6 5	108.50	110.50
156	2 6 1	113.50	113.50
157	2 6 2	111.50	112.00
158	2 6 3	114.50	116.50
159	2 6 4	120.00	121.50
160	2 6 5	111.00	114.00
161	3 6 1	091.50	092.50
162	3 6 2	093.00	095.00
163	3 6 3	094.50	097.50
164	3 6 4	101.50	103.50
165	3 6 5	095.00	099.00
166	4 6 1	104.00	105.50
167	4 6 2	105.50	104.00
168	4 6 3	101.50	104.50
169	4 6 4	107.00	105.50
170	4 6 5	104.00	107.00
171	5 6 1	101.50	102.00
172	5 6 2	100.50	101.50
173	5 6 3	102.50	104.50
174	5 6 4	104.00	105.00
175	5 6 5	102.50	103.50
176	6 6 1	105.50	105.50
177	6 6 2	104.50	104.50
178	6 6 3	105.50	105.00
179	6 6 4	107.50	104.50
180	6 6 5	102.50	107.00

#	SAB	C_1	C_2
181	1 7 1	094.50	095.00
182	1 7 2	099.50	096.00
183	1 7 3	091.50	092.00
184	1 7 4	097.50	097.50
185	1 7 5	093.00	098.00
186	2 7 1	104.50	106.00
187	2 7 2	104.50	105.50
188	2 7 3	104.50	103.50
189	2 7 4	107.00	104.50
190	2 7 5	104.50	104.50
191	3 7 1	116.50	116.50
192	3 7 2	114.00	113.00
193	3 7 3	113.50	114.00
194	3 7 4	120.50	122.50
195	3 7 5	116.50	116.00
196	4 7 1	104.50	104.50
197	4 7 2	105.50	107.00
198	4 7 3	105.50	105.50
199	4 7 4	108.00	107.50
200	4 7 5	105.50	107.00
201	5 7 1	104.50	104.50
202	5 7 2	110.50	110.00
203	5 7 3	103.50	103.50
204	5 7 4	107.50	108.50
205	5 7 5	110.00	110.50
206	6 7 1	105.00	105.00
207	6 7 2	104.00	106.00
208	6 7 3	103.50	103.00
209	6 7 4	104.50	104.00
210	6 7 5	104.00	103.50
211	SENDFILE		

END OF FILE

Symbols

- # Number
- S Subject
- A Treatment
- B Week
- C Session

Treatment Conditions

- A_1 3S training (n = 6)
- A_2 2S training (n = 6)
- A_3 SS training (n = 6)
- A_4 ES training (n = 6)
- A_5 BS training (n = 6)
- A_6 CS training (n = 6)
- A_7 Control (n = 6)

Week in Experiment

- B_1 pre-training
- B_2 two-week training
- B_3 four-week training
- B_4 six-week training
- B_5 two-week retention

Session

- C_1 pre-session training
- C_2 post-session training

TABLE C

RAW SCORES (IN DEGREES) OF SHOULDER JOINT FLEXIBILITY FOR EACH SUBJECT

#	S	A	B	C	C_1	C_2
1	1	1	1		155.00	155.00
2	1	1	2		162.00	166.00
3	1	1	3		164.00	166.00
4	1	1	4		166.00	166.00
5	1	1	5		168.00	168.00
6	2	1	1		142.00	146.00
7	2	1	2		160.00	166.00
8	2	1	3		160.00	162.00
9	2	1	4		167.00	166.00
10	2	1	5		165.00	167.00
11	3	1	1		163.00	167.00
12	3	1	2		167.00	160.00
13	3	1	3		167.00	170.00
14	3	1	4		176.00	176.00
15	3	1	5		160.00	167.00
16	4	1	1		148.00	166.00
17	4	1	2		160.00	162.00
18	4	1	3		164.00	160.00
19	4	1	4		174.00	176.00
20	4	1	5		163.00	160.00
21	5	1	1		121.00	126.00
22	5	1	2		166.00	166.00
23	5	1	3		163.00	174.00
24	5	1	4		164.00	164.00
25	5	1	5		164.00	163.00
26	6	1	1		148.00	166.00
27	6	1	2		166.00	171.00
28	6	1	3		174.00	174.00
29	6	1	4		168.00	170.00
30	6	1	5		164.00	168.00
31	1	2	1		148.00	148.00
32	1	2	2		166.00	166.00
33	1	2	3		164.00	168.00
34	1	2	4		166.00	165.00
35	1	2	5		163.00	161.00
36	2	2	1		141.00	146.00
37	2	2	2		161.00	161.00
38	2	2	3		160.00	161.00
39	2	2	4		165.00	166.00
40	2	2	5		160.00	161.00
41	3	2	1		157.00	163.00
42	3	2	2		163.00	163.00
43	3	2	3		160.00	163.00
44	3	2	4		164.00	166.00
45	3	2	5		161.00	160.00
46	4	2	1		150.00	148.00
47	4	2	2		166.00	166.00
48	4	2	3		166.00	163.00
49	4	2	4		176.00	174.00
50	4	2	5		173.00	166.00
51	5	2	1		160.00	164.00
52	5	2	2		164.00	166.00
53	5	2	3		176.00	174.00
54	5	2	4		173.00	171.00
55	5	2	5		173.00	171.00
56	6	2	1		168.00	166.00
57	6	2	2		168.00	171.00
58	6	2	3		173.00	174.00
59	6	2	4		170.00	169.00
60	6	2	5		171.00	175.00
61	1	3	1		152.00	154.00
62	1	3	2		168.00	166.00
63	1	3	3		165.00	160.00
64	1	3	4		174.00	170.00
65	1	3	5		166.00	166.00
66	2	3	1		136.00	136.00
67	2	3	2		162.00	161.00
68	2	3	3		160.00	161.00
69	2	3	4		170.00	170.00
70	2	3	5		168.00	160.00
71	3	3	1		148.00	148.00
72	3	3	2		160.00	162.00
73	3	3	3		160.00	167.00
74	3	3	4		167.00	168.00
75	3	3	5		166.00	166.00
76	4	3	1		162.00	160.00
77	4	3	2		167.00	167.00
78	4	3	3		164.00	162.00
79	4	3	4		170.00	168.00
80	4	3	5		160.00	162.00
81	5	3	1		161.00	164.00
82	5	3	2		167.00	164.00
83	5	3	3		166.00	166.00
84	5	3	4		174.00	174.00
85	5	3	5		166.00	167.00
86	6	3	1		161.00	161.00
87	6	3	2		174.00	174.00
88	6	3	3		160.00	170.00
89	6	3	4		174.00	168.00
90	6	3	5		168.00	168.00
91	1	4	1		166.00	168.00
92	1	4	2		164.00	162.00
93	1	4	3		164.00	162.00
94	1	4	4		174.00	170.00
95	1	4	5		163.00	167.00
96	2	4	1		138.00	140.00
97	2	4	2		167.00	166.00
98	2	4	3		162.00	164.00
99	2	4	4		174.00	174.00
100	2	4	5		162.00	162.00
101	3	4	1		166.00	160.00
102	3	4	2		162.00	164.00
103	3	4	3		166.00	166.00
104	3	4	4		162.00	162.00
105	3	4	5		168.00	174.00
106	4	4	1		168.00	168.00
107	4	4	2		177.00	180.00
108	4	4	3		170.00	180.00
109	4	4	4		171.00	171.00
110	4	4	5		171.00	172.00
111	5	4	1		167.00	166.00
112	5	4	2		167.00	172.00
113	5	4	3		172.00	172.00
114	5	4	4		172.00	172.00
115	5	4	5		164.00	166.00
116	6	4	1		138.00	146.00
117	6	4	2		166.00	166.00
118	6	4	3		162.00	167.00
119	6	4	4		170.00	168.00
120	6	4	5		162.00	168.00

Symbols

- # Number
- S Subject
- A Treatment
- B Week
- C Session

Treatment Conditions

- A_1 3S training (n = 6)
- A_2 2S training (n = 6)
- A_3 SS training (n = 6)
- A_4 ES training (n = 6)
- A_5 BS training (n = 6)
- A_6 CS training (n = 6)
- A_7 Control (n = 6)

Week in Experiment

- B_1 pre-training
- B_2 two-week training
- B_3 four-week training
- B_4 six-week training
- B_5 two-week retention

Session

- C_1 pre-session training
- C_2 post-session training

University of Alberta

TABLE D

RAW SCORES (IN DEGREES) OF SHOULDER JOINT FLEXIBILITY FOR EACH SUBJECT

#	SAB	C_1	C_2
121	1 6 1	166.50	152.00
122	1 6 2	187.50	188.00
123	1 6 3	161.00	162.50
124	1 6 4	176.50	170.00
125	1 6 5	161.50	160.50
126	2 6 1	164.50	164.00
127	2 6 2	183.00	180.00
128	2 6 3	187.00	185.00
129	2 6 4	174.00	173.50
130	2 6 5	186.00	183.00
131	3 6 1	146.00	146.00
132	3 6 2	158.00	158.50
133	3 6 3	178.00	174.50
134	3 6 4	171.50	172.00
135	3 6 5	189.00	188.00
136	4 6 1	158.50	152.00
137	4 6 2	182.50	184.00
138	4 6 3	181.50	180.00
139	4 6 4	180.00	184.50
140	4 6 5	172.50	168.00
141	5 6 1	181.00	185.50
142	5 6 2	180.00	185.50
143	5 6 3	178.00	180.00
144	5 6 4	178.50	178.50
145	5 6 5	172.50	172.50
146	6 6 1	135.50	138.00
147	6 6 2	180.50	182.00
148	6 6 3	152.50	153.00
149	6 6 4	153.50	151.00
150	6 6 5	148.50	150.00
151	1 7 1	145.00	144.50
152	1 7 2	158.00	158.50
153	1 7 3	198.00	188.50
154	1 7 4	183.50	185.50
155	1 7 5	182.50	182.00
156	2 7 1	182.00	180.50
157	2 7 2	182.00	182.50
158	2 7 3	188.50	186.00
159	2 7 4	174.00	172.00
160	2 7 5	171.50	170.00
161	3 7 1	140.00	141.00
162	3 7 2	183.50	184.00
163	3 7 3	187.50	189.50
164	3 7 4	176.50	177.00
165	3 7 5	183.50	180.50
166	4 7 1	138.00	138.50
167	4 7 2	180.00	181.00
168	4 7 3	141.00	140.00
169	4 7 4	170.50	170.50
170	4 7 5	158.50	162.50
171	5 7 1	181.00	181.00
172	5 7 2	187.50	189.50
173	5 7 3	170.00	172.00
174	5 7 4	172.50	173.00
175	5 7 5	182.50	180.00
176	6 7 1	158.50	160.00
177	6 7 2	171.00	174.50
178	6 7 3	180.50	180.50
179	6 7 4	165.00	163.00
180	6 7 5	170.50	170.00

#	SAB	C_1	C_2
181	1 7 1	143.50	142.00
182	1 7 2	182.00	183.50
183	1 7 3	198.00	198.50
184	1 7 4	174.50	172.50
185	1 7 5	182.50	198.50
186	2 7 1	144.50	145.00
187	2 7 2	198.50	186.50
188	2 7 3	148.00	148.00
189	2 7 4	151.50	151.00
190	2 7 5	141.50	143.00
191	3 7 1	134.00	134.00
192	3 7 2	144.00	148.00
193	3 7 3	134.50	137.00
194	3 7 4	158.00	160.50
195	3 7 5	160.00	158.50
196	4 7 1	158.50	158.00
197	4 7 2	158.50	171.50
198	4 7 3	177.50	180.00
199	4 7 4	172.00	171.50
200	4 7 5	171.50	172.50
201	5 7 1	138.00	138.50
202	5 7 2	148.50	148.50
203	5 7 3	148.50	148.00
204	5 7 4	185.00	184.50
205	5 7 5	148.50	182.50
206	6 7 1	144.50	143.50
207	6 7 2	158.50	180.50
208	6 7 3	184.50	183.50
209	6 7 4	157.50	158.50
210	6 7 5	145.50	144.00
211	SENDFILE		

END OF FILE

Symbols

- # Number
- S Subject
- A Treatment
- B Week
- C Session

Week in Experiment

- B_1 pre-training
- B_2 two-week training
- B_3 four-week training
- B_4 six-week training
- B_5 two-week retention

Treatment Conditions

- A_1 3S training (n = 6)
- A_2 2S training (n = 6)
- A_3 SS training (n = 6)
- A_4 ES training (n = 6)
- A_5 BS training (n = 6)
- A_6 CS training (n = 6)
- A_7 Control (n = 6)

Session

- C_1 pre-session training
- C_2 post-session training

APPENDIX E

TABLES OF MEANS

TABLE E

TREATMENT(A) BY WEEK(B) BY SESSION(C) SUMMARY TABLE
OF CELL MEANS FOR ANKLE DATA

	B_1		B_2		B_3		B_4		B_5	
	C_1	C_2	C_1	C_2	C_1	C_2	C_1	C_2	C_1	C_2
A_1	107.50	112.17	106.17	109.58	107.67	111.50	113.42	115.67	109.00	113.25
A_2	104.58	107.50	104.33	107.75	105.00	107.58	110.67	113.67	105.92	108.92
A_3	105.33	104.92	104.83	104.58	105.42	104.50	108.92	108.75	106.83	105.92
A_4	104.75	106.42	104.08	107.25	105.00	107.42	107.92	111.25	105.25	108.83
A_5	104.67	106.42	106.67	106.33	107.08	107.42	110.67	111.25	105.00	105.58
A_6	104.42	105.08	104.17	106.42	104.50	105.92	109.08	110.42	104.08	106.83
A_7	104.92	105.08	105.67	105.75	103.67	103.58	107.17	107.75	106.08	106.08
Average Means	105.17	106.79	105.13	106.81	105.48	106.85	109.69	111.25	106.02	107.92

TABLE F

TREATMENT(A) BY WEEK(B) SUMTARY TABLE
OF CELL MEANS FOR ANKLE DATA

	B_1	B_2	B_3	B_4	B_5	Average Means
A_1	109.83	107.88	109.58	114.54	111.13	110.59
A_2	106.04	106.04	106.29	112.17	107.42	107.59
A_3	105.13	104.71	104.96	108.83	106.38	106.00
A_4	105.58	105.67	106.21	109.58	107.04	106.82
A_5	105.54	106.50	107.25	110.96	105.29	107.11
A_6	104.75	105.29	105.21	109.75	105.46	106.09
A_7	105.00	105.71	103.63	107.46	106.08	105.58
Average Means	105.98	105.97	106.16	110.47	106.97	107.11

TABLE G

TREATMENT(A) BY SESSION(C) SUMMARY TABLE
OF CELL MEANS FOR ANKLE DATA

	C_1	C_2	Average Means
A_1	108.75	112.43	110.59
A_2	106.10	109.08	107.59
A_3	106.27	105.73	106.00
A_4	105.40	108.23	106.82
A_5	106.82	107.40	107.11
A_6	105.25	106.93	106.09
A_7	105.50	105.65	105.58
Average Means	106.29	107.92	107.11

TABLE H

WEEK(B) BY SESSION(C) SUMMARY TABLE
OF CELL MEANS FOR ANKLE DATA

	C_1	C_2	Average Means
B_1	105.17	106.79	105.98
B_2	105.13	106.81	105.97
B_3	105.48	106.85	106.16
B_4	109.69	111.25	110.47
B_5	106.02	107.92	106.97
Average Means	106.29	107.92	107.11

TABLE I

TREATMENT(A) BY WEEK(B) BY SESSION(C) SUMMARY TABLE
OF CELL MEANS FOR SHOULDER DATA

	B_1		B_2		B_3		B_4		B_5	
	C_1	C_2	C_1	C_2	C_1	C_2	C_1	C_2	C_1	C_2
A_1	144.08	151.00	160.58	163.08	164.08	166.75	172.17	175.17	164.25	167.33
A_2	150.50	152.67	164.75	165.75	169.08	170.92	175.50	175.42	168.58	167.58
A_3	150.33	150.58	162.42	161.17	163.92	162.92	171.92	169.92	162.75	161.75
A_4	150.67	153.50	164.00	165.50	165.92	168.83	175.25	174.67	160.58	163.17
A_5	148.83	149.25	161.25	161.58	166.00	166.00	173.00	171.25	163.67	163.33
A_6	152.25	152.25	161.75	163.00	162.50	164.58	174.67	174.50	163.17	163.50
A_7	143.58	143.33	154.33	155.58	153.83	154.17	161.58	161.58	155.08	155.17
Average Means	148.61	150.37	161.29	162.24	163.62	164.88	172.01	171.79	162.58	163.12

TABLE J

TREATMENT(A) BY WEEK(B) SUMMARY TABLE
OF CELL MEANS FOR SHOULDER DATA

	B_1	B_2	B_3	B_4	B_5	Average Means
A_1	147.54	161.83	165.42	173.67	165.79	162.85
A_2	151.58	165.25	170.00	175.46	168.08	166.08
A_3	150.46	161.79	163.42	170.92	162.25	161.77
A_4	152.08	164.75	167.38	174.96	161.88	164.21
A_5	149.04	161.42	166.00	172.13	163.50	162.42
A_6	152.25	162.38	163.54	174.58	163.33	163.22
A_7	143.46	154.96	154.00	161.58	155.13	153.83
Average Means	149.49	161.77	164.25	171.89	162.85	162.05

TABLE K

TREATMENT(A) BY SESSION(C) SUMMARY TABLE
OF CELL MEANS FOR SHOULDER DATA

	C_1	C_2	Average Means
A_1	161.03	164.67	162.85
A_2	165.68	166.47	166.08
A_3	162.27	161.27	161.77
A_4	163.28	165.13	164.21
A_5	162.55	162.28	162.42
A_6	162.87	163.57	163.22
A_7	153.68	153.97	153.83
Average Means	161.62	162.48	162.05

TABLE L

WEEK(B) BY SESSION(C) SUMMARY TABLE
OF CELL MEANS FOR SHOULDER DATA

	C_1	C_2	Average Means
B_1	148.61	150.37	149.49
B_2	161.29	162.24	161.77
B_3	163.62	164.88	164.25
B_4	172.01	171.79	171.89
B_5	162.58	163.12	162.85
Average Means	161.62	162.48	162.05